



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 07:04 PM EDT

PDB ID : 4QPD
Title : Crystal structure of the hydrolase domain of 10-formyltetrahydrofolate dehydrogenase (wild-type) complex with tetrahydrofolate
Authors : Lin, C.C.; Chen, C.J.; Fu, T.F.; Chuankhayan, P.; Kao, T.T.; Chang, W.N.
Deposited on : 2014-06-23
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	<div> <div>10%</div> <div>90%</div> <div>6% . .</div> </div>
1	B	318	<div> <div>9%</div> <div>87%</div> <div>10% .</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 10-formyltetrahydrofolate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	S	0	0	0
			2415	1549	415	442	9			
1	B	308	Total	C	N	O	S	0	0	0
			2415	1549	415	442	9			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	312	GLU	-	expression tag	UNP E3NZ06
A	313	HIS	-	expression tag	UNP E3NZ06
A	314	HIS	-	expression tag	UNP E3NZ06
A	315	HIS	-	expression tag	UNP E3NZ06
A	316	HIS	-	expression tag	UNP E3NZ06
A	317	HIS	-	expression tag	UNP E3NZ06
A	318	HIS	-	expression tag	UNP E3NZ06
B	312	GLU	-	expression tag	UNP E3NZ06
B	313	HIS	-	expression tag	UNP E3NZ06
B	314	HIS	-	expression tag	UNP E3NZ06
B	315	HIS	-	expression tag	UNP E3NZ06
B	316	HIS	-	expression tag	UNP E3NZ06
B	317	HIS	-	expression tag	UNP E3NZ06
B	318	HIS	-	expression tag	UNP E3NZ06

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is (6S)-5,6,7,8-TETRAHYDROFOLATE (three-letter code: THG) (formula: $C_{19}H_{23}N_7O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			32	19	7	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			32	19	7	6		
3	B	1	Total	C	N	O	0	0
			32	19	7	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	205	Total	O	0	0
			205	205		
4	B	227	Total	O	0	0
			227	227		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 10-formyltetrahydrofolate dehydrogenase



- Molecule 1: 10-formyltetrahydrofolate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.50Å 100.42Å 122.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.37 – 2.10 29.37 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.7 (29.37-2.10) 98.4 (29.37-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.185 , 0.233 0.187 , 0.235	Depositor DCC
R_{free} test set	1987 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5372	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3371e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: THG, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/2475	0.46	1/3348 (0.0%)
1	B	0.29	0/2475	0.46	0/3348
All	All	0.29	0/4950	0.46	1/6696 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	288	LEU	CA-CB-CG	6.86	131.09	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	0	2411	10	0
1	B	2415	0	2411	18	0
2	A	7	0	10	0	0
2	B	7	0	10	0	0
3	A	32	0	21	1	0
3	B	64	0	42	4	0
4	A	205	0	0	0	0
4	B	227	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5372	0	4905	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:258:ASN:HA	3:B:402:THG:HCG2	1.68	0.76
1:A:255:PHE:N	1:A:256:SER:HA	2.13	0.63
1:B:116:ALA:HB3	1:B:203:ILE:HG12	1.82	0.61
1:A:108:SER:HB3	1:A:118:ALA:HB1	1.81	0.60
1:B:205:LYS:NZ	4:B:642:HOH:O	2.36	0.58
1:A:32:ILE:HD12	1:A:35:LYS:HE3	1.84	0.58
1:B:299:LYS:NZ	4:B:693:HOH:O	2.39	0.55
1:A:228:ASP:OD2	1:A:229:ARG:HD2	2.10	0.51
1:B:73:TYR:O	1:B:76:VAL:HG22	2.12	0.50
1:B:258:ASN:CA	3:B:402:THG:HCG2	2.40	0.49
1:B:136:TRP:O	1:B:144:GLY:HA3	2.12	0.48
1:A:203:ILE:HG21	3:A:402:THG:HCG1	1.95	0.47
1:B:164:TYR:HA	1:B:168:LEU:HB2	1.97	0.47
1:B:255:PHE:CE2	1:B:274:THR:HB	2.51	0.45
1:B:285:GLY:HA2	3:B:402:THG:N1	2.32	0.45
1:B:108:SER:HB3	1:B:118:ALA:HB1	1.99	0.44
1:A:260:GLN:HA	1:A:261:PRO:HD3	1.84	0.44
1:B:135:PHE:CZ	1:B:142:ASP:HA	2.53	0.43
3:B:402:THG:HCB2	3:B:402:THG:O11	2.17	0.43
1:B:61:LEU:HD23	1:B:61:LEU:HA	1.88	0.43
1:A:136:TRP:O	1:A:144:GLY:HA3	2.18	0.43
1:B:92:MET:HA	1:B:95:ILE:HB	2.01	0.43
1:B:255:PHE:HE1	1:B:276:ASN:HD22	1.67	0.42
1:A:38:LYS:HE3	1:A:38:LYS:HB2	1.71	0.42
1:B:40:ASP:HA	1:B:41:PRO:HD3	1.91	0.42
1:B:31:THR:HG21	1:B:43:ALA:HB2	2.00	0.42
1:B:228:ASP:HA	1:B:229:ARG:HA	1.74	0.42
1:A:81:ASN:HB2	1:A:102:SER:OG	2.20	0.42
1:A:164:TYR:HA	1:A:168:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/318 (96%)	292 (95%)	13 (4%)	1 (0%)	41	41
1	B	306/318 (96%)	296 (97%)	10 (3%)	0	100	100
All	All	612/636 (96%)	588 (96%)	23 (4%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/266 (97%)	254 (98%)	4 (2%)	62	69
1	B	258/266 (97%)	254 (98%)	4 (2%)	62	69
All	All	516/532 (97%)	508 (98%)	8 (2%)	62	69

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	ASP
1	A	256	SER
1	A	275	LYS
1	A	288	LEU
1	B	35	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	47	GLU
1	B	84	PRO
1	B	252	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	THG	B	403	-	32,34,34	1.75	8 (25%)	39,47,47	1.66	8 (20%)
3	THG	B	402	-	32,34,34	1.71	8 (25%)	39,47,47	2.25	11 (28%)
2	PEG	A	401	-	6,6,6	0.70	0	5,5,5	0.20	0
2	PEG	B	401	-	6,6,6	0.70	0	5,5,5	0.24	0
3	THG	A	402	-	32,34,34	1.71	8 (25%)	39,47,47	1.60	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	THG	B	403	-	-	4/22/31/31	0/3/3/3
3	THG	B	402	-	-	6/22/31/31	0/3/3/3
2	PEG	A	401	-	-	1/4/4/4	-
2	PEG	B	401	-	-	1/4/4/4	-
3	THG	A	402	-	-	1/22/31/31	0/3/3/3

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	THG	C2-N2	3.59	1.41	1.33
3	A	402	THG	C4A-C4	-3.51	1.36	1.41
3	B	403	THG	C2-N2	3.49	1.40	1.33
3	B	402	THG	C2-N2	3.32	1.40	1.33
3	B	403	THG	C4A-C4	-3.30	1.37	1.41
3	B	402	THG	C11-N	3.28	1.41	1.34
3	B	402	THG	C7-N8	-3.19	1.39	1.44
3	B	403	THG	C4'-N10	3.16	1.47	1.38
3	B	402	THG	C4A-N5	3.11	1.44	1.38
3	B	402	THG	C4'-N10	3.05	1.47	1.38
3	A	402	THG	C4'-N10	3.05	1.47	1.38
3	B	403	THG	C1'-C11	3.02	1.56	1.50
3	B	403	THG	C4A-N5	3.01	1.44	1.38
3	A	402	THG	C4A-N5	2.98	1.44	1.38
3	B	402	THG	C4A-C4	-2.96	1.37	1.41
3	A	402	THG	C1'-C11	2.85	1.56	1.50
3	B	403	THG	C11-N	2.71	1.40	1.34
3	B	403	THG	C7-N8	-2.71	1.40	1.44
3	B	402	THG	C1'-C11	2.64	1.55	1.50
3	A	402	THG	C7-N8	-2.64	1.40	1.44
3	A	402	THG	C11-N	2.55	1.39	1.34
3	B	402	THG	C5'-C6'	2.11	1.42	1.38
3	A	402	THG	C3'-C4'	2.03	1.42	1.39
3	B	403	THG	C2'-C1'	2.02	1.42	1.39

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	THG	C4-C4A-N5	7.16	125.13	119.12
3	B	403	THG	C4-C4A-N5	5.49	123.72	119.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	THG	C4-C4A-N5	5.34	123.60	119.12
3	B	402	THG	CG-CB-CA	-5.34	103.18	113.16
3	B	402	THG	C6-C7-N8	-4.65	101.95	110.68
3	B	402	THG	CA-N-C11	4.02	131.44	121.60
3	B	402	THG	C4A-N5-C6	-3.93	111.15	121.48
3	A	402	THG	C8A-C4A-C4	3.38	117.57	114.57
3	B	403	THG	C8A-C4A-C4	3.29	117.49	114.57
3	A	402	THG	C4A-N5-C6	-2.92	113.81	121.48
3	B	402	THG	C8A-C4A-C4	2.88	117.13	114.57
3	B	403	THG	C4A-N5-C6	-2.83	114.02	121.48
3	B	403	THG	C2-N1-C8A	2.68	120.55	114.54
3	B	402	THG	CB-CA-N	2.44	115.81	110.88
3	A	402	THG	C6-C7-N8	-2.42	106.13	110.68
3	B	402	THG	C2-N1-C8A	2.42	119.96	114.54
3	B	403	THG	OX2-C-CA	2.40	121.37	113.40
3	A	402	THG	C2-N1-C8A	2.35	119.80	114.54
3	B	403	THG	CG-CB-CA	-2.33	108.80	113.16
3	B	403	THG	C6-C7-N8	-2.29	106.38	110.68
3	A	402	THG	OX2-C-CA	2.16	120.60	113.40
3	B	402	THG	C1'-C11-N	-2.08	113.08	117.06
3	B	402	THG	N2-C2-N3	2.05	120.44	117.25
3	B	403	THG	C1'-C11-N	2.04	120.97	117.06
3	B	402	THG	C9-C6-N5	2.03	113.20	108.83

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	402	THG	CB-CA-N-C11
3	B	402	THG	N-CA-CB-CG
3	B	403	THG	N-CA-CB-CG
3	B	402	THG	C-CA-CB-CG
3	B	403	THG	C-CA-CB-CG
2	A	401	PEG	O1-C1-C2-O2
2	B	401	PEG	O2-C3-C4-O4
3	A	402	THG	C-CA-CB-CG
3	B	403	THG	OE1-CD-CG-CB
3	B	403	THG	OE2-CD-CG-CB
3	B	402	THG	OE1-CD-CG-CB
3	B	402	THG	C6-C9-N10-C4'
3	B	402	THG	OE2-CD-CG-CB



There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	THG	4	0
3	A	402	THG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/318 (96%)	0.59	31 (10%) 7 9	14, 30, 66, 165	0
1	B	308/318 (96%)	0.53	28 (9%) 9 12	13, 27, 67, 172	0
All	All	616/636 (96%)	0.56	59 (9%) 8 10	13, 28, 67, 172	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	255	PHE	14.5
1	B	255	PHE	11.5
1	B	254	HIS	10.1
1	A	297	ASP	9.1
1	A	254	HIS	8.0
1	A	39	VAL	7.8
1	B	257	SER	7.7
1	B	252	ASN	7.7
1	B	253	ASP	7.0
1	B	256	SER	7.0
1	B	297	ASP	6.4
1	A	253	ASP	6.2
1	A	252	ASN	6.1
1	B	37	GLY	5.7
1	A	256	SER	5.2
1	A	258	ASN	5.1
1	A	38	LYS	5.0
1	A	36	ASP	5.0
1	A	257	SER	4.9
1	A	296	GLU	4.8
1	B	36	ASP	4.5
1	A	35	LYS	4.3
1	A	43	ALA	4.1
1	B	35	LYS	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	38	LYS	3.6
1	A	37	GLY	3.6
1	B	258	ASN	3.2
1	B	299	LYS	3.1
1	B	39	VAL	3.0
1	A	32	ILE	2.9
1	A	207	GLU	2.9
1	B	34	ASP	2.7
1	B	207	GLU	2.7
1	A	54	PHE	2.7
1	B	307	PHE	2.7
1	A	134	VAL	2.6
1	A	290	VAL	2.6
1	A	104	ILE	2.5
1	B	195	GLU	2.5
1	B	238	ASP	2.4
1	A	224	ILE	2.4
1	A	299	LYS	2.3
1	A	308	LYS	2.3
1	B	62	LYS	2.3
1	B	83	LEU	2.3
1	B	280	LEU	2.3
1	A	280	LEU	2.3
1	A	62	LYS	2.3
1	A	288	LEU	2.2
1	B	300	MET	2.2
1	A	250	LEU	2.2
1	B	68	GLU	2.2
1	B	288	LEU	2.1
1	B	296	GLU	2.1
1	A	33	PRO	2.1
1	A	46	ALA	2.1
1	B	308	LYS	2.1
1	A	85	PHE	2.1
1	B	40	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PEG	B	401	7/7	0.64	0.27	53,55,57,57	0
2	PEG	A	401	7/7	0.67	0.24	51,51,52,54	0
3	THG	B	402	32/32	0.88	0.27	18,49,84,85	0
3	THG	A	402	32/32	0.90	0.17	17,27,52,55	0
3	THG	B	403	32/32	0.91	0.18	16,23,62,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around THG A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around THG B 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.