



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2025 – 02:57 PM EDT

PDB ID : 2RII / pdb\_00002rii  
Title : Crystal Structure of Human Peroxiredoxin I in complex with Sulfiredoxin  
Authors : Jonsson, T.J.; Johnson, L.C.; Lowther, W.T.  
Deposited on : 2007-10-11  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	
1	B	199	
2	X	110	
2	Y	110	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4485 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxiredoxin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	184	Total	C	N	O	S	0	0	0
			1435	926	240	266	3			
1	B	185	Total	C	N	O	S	0	0	0
			1443	930	241	269	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	71	SER	CYS	engineered mutation	UNP Q06830
A	83	GLU	CYS	engineered mutation	UNP Q06830
A	173	SER	CYS	engineered mutation	UNP Q06830
B	71	SER	CYS	engineered mutation	UNP Q06830
B	83	GLU	CYS	engineered mutation	UNP Q06830
B	173	SER	CYS	engineered mutation	UNP Q06830

- Molecule 2 is a protein called Sulfiredoxin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	106	Total	C	N	O	S	0	0	0
			820	523	142	154	1			
2	Y	99	Total	C	N	O	S	0	0	0
			767	491	130	145	1			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	28	GLY	-	expression tag	UNP Q9BYN0
X	29	HIS	-	expression tag	UNP Q9BYN0
X	30	PRO	-	expression tag	UNP Q9BYN0
X	31	MET	-	expression tag	UNP Q9BYN0
Y	28	GLY	-	expression tag	UNP Q9BYN0
Y	29	HIS	-	expression tag	UNP Q9BYN0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Y	30	PRO	-	expression tag	UNP Q9BYN0
Y	31	MET	-	expression tag	UNP Q9BYN0

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	X	1	Total	O	P	0	0
			5	4	1		
3	Y	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	O	0	0
			2	2		
4	B	4	Total	O	0	0
			4	4		
4	X	3	Total	O	0	0
			3	3		
4	Y	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Peroxiredoxin-1




#### • Molecule 1: Peroxiredoxin-1



#### • Molecule 2: Sulfiredoxin-1



#### • Molecule 2: Sulfiredoxin-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.92Å 85.05Å 130.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.52 – 2.60 43.52 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.52-2.60) 99.9 (43.52-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.59 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.239 , 0.308 0.235 , 0.306	Depositor DCC
$R_{free}$ test set	995 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.0	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 13.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4485	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	0/1473	1.02	2/1996 (0.1%)
1	B	0.82	0/1481	1.05	2/2007 (0.1%)
2	X	0.91	1/840 (0.1%)	1.07	4/1147 (0.3%)
2	Y	0.70	0/786	0.93	0/1075
All	All	0.80	1/4580 (0.0%)	1.02	8/6225 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	X	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	X	125	LEU	C-O	-6.72	1.16	1.24

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	X	125	LEU	O-C-N	-10.85	110.37	122.09
1	B	147	LEU	N-CA-C	5.84	122.71	109.81
1	A	149	VAL	N-CA-C	5.83	117.40	108.71
2	X	125	LEU	CA-C-O	5.54	126.41	120.70
2	X	53	LEU	CA-C-N	5.34	126.52	119.84
2	X	53	LEU	C-N-CA	5.34	126.52	119.84
1	B	149	VAL	N-CA-C	5.18	115.52	107.28
1	A	171	GLU	N-CA-C	-5.12	106.88	112.72

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	X	125	LEU	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1435	0	1421	57	0
1	B	1443	0	1425	34	0
2	X	820	0	821	24	0
2	Y	767	0	766	23	0
3	X	5	0	0	1	0
3	Y	5	0	0	1	0
4	A	2	0	0	0	0
4	B	4	0	0	0	0
4	X	3	0	0	0	0
4	Y	1	0	0	0	0
All	All	4485	0	4433	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:141:GLN:HE22	1:B:143:THR:HG23	0.99	1.15
1:A:140:ARG:HG2	1:A:140:ARG:HH11	1.13	1.12
1:A:181:SER:HB3	1:A:182:ASP:HA	1.34	1.05
1:B:141:GLN:NE2	1:B:143:THR:HG23	1.73	1.02
1:A:141:GLN:HE22	1:A:143:THR:CG2	1.74	0.99
1:A:140:ARG:HH11	1:A:140:ARG:CG	1.79	0.95
2:Y:45:PRO:O	2:Y:48:VAL:HG12	1.67	0.93
1:A:141:GLN:HE22	1:A:143:THR:HG23	1.31	0.93
1:A:134:ASP:HB3	1:A:140:ARG:HD2	1.48	0.91
1:A:181:SER:HA	1:A:183:THR:H	1.40	0.85

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ARG:HG2	1:A:140:ARG:NH1	1.91	0.83
1:A:175:ALA:CB	2:Y:48:VAL:HG23	2.11	0.81
1:A:141:GLN:NE2	1:A:143:THR:HG23	1.96	0.79
1:A:175:ALA:HB1	2:Y:48:VAL:HG23	1.65	0.77
1:A:141:GLN:HE22	1:A:143:THR:HG22	1.54	0.72
1:A:141:GLN:NE2	1:A:143:THR:CG2	2.50	0.71
1:B:155:GLU:OE1	1:B:158:ARG:NH1	2.24	0.70
1:A:134:ASP:HB3	1:A:140:ARG:CD	2.22	0.69
1:B:178:LYS:HD2	2:X:91:ASP:OD1	1.93	0.69
1:A:150:GLY:O	1:B:158:ARG:NH2	2.28	0.67
1:A:47:ASP:O	1:A:87:TRP:CE2	2.47	0.67
2:Y:58:ASP:HB3	2:Y:61:LYS:HD2	1.75	0.67
1:B:155:GLU:CD	1:B:158:ARG:NH1	2.54	0.65
1:B:179:PRO:HG3	2:X:117:LEU:HD13	1.81	0.63
1:B:183:THR:HG22	2:X:42:HIS:ND1	2.15	0.62
1:B:100:MET:HA	1:B:100:MET:HE2	1.82	0.61
2:Y:97:GLY:O	2:Y:99:CYS:N	2.33	0.61
1:A:134:ASP:CB	1:A:140:ARG:HD2	2.26	0.61
1:A:15:PHE:CE2	1:A:31:LEU:HB2	2.37	0.60
1:B:155:GLU:CD	1:B:158:ARG:HH11	2.10	0.58
1:A:55:GLU:OE2	1:A:151:ARG:HD3	2.04	0.58
2:X:44:VAL:CG1	2:X:115:ALA:HB3	2.34	0.57
1:A:166:THR:HG22	2:Y:54:PRO:HD3	1.85	0.57
1:B:174:PRO:HD3	2:X:92:TYR:OH	2.05	0.57
2:X:38:ILE:O	2:X:39:ALA:HB2	2.05	0.56
1:A:50:PHE:CD2	2:X:82:LEU:HD11	2.41	0.56
1:A:123:GLU:HB2	1:A:125:ILE:HG12	1.87	0.56
1:B:128:ARG:NH1	1:B:147:LEU:O	2.38	0.56
2:Y:85:LYS:HA	2:Y:90:GLY:O	2.07	0.55
2:Y:44:VAL:HG23	2:Y:48:VAL:CG1	2.37	0.55
2:X:51:ARG:HD2	2:X:102:TYR:CD2	2.42	0.54
1:A:73:VAL:HB	1:A:102:ILE:HG21	1.89	0.54
1:A:21:MET:HB3	1:A:22:PRO:HD2	1.90	0.54
1:A:140:ARG:CG	1:A:140:ARG:NH1	2.53	0.54
1:A:181:SER:HB3	1:A:182:ASP:CA	2.24	0.54
2:X:37:ARG:HD2	2:X:37:ARG:N	2.23	0.54
2:X:37:ARG:HB3	2:X:38:ILE:HA	1.91	0.53
1:A:3:SER:HA	1:B:3:SER:N	2.25	0.51
1:A:32:SER:O	1:A:35:LYS:HB2	2.10	0.51
1:A:50:PHE:HD2	2:X:82:LEU:HD11	1.76	0.51
1:B:140:ARG:HB3	1:B:163:PHE:CE2	2.45	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:ARG:HD2	1:A:65:GLU:OE2	2.10	0.50
1:B:51:VAL:HG23	2:Y:128:TYR:CE2	2.45	0.50
2:Y:97:GLY:O	2:Y:98:GLY:C	2.55	0.50
1:A:158:ARG:NH2	1:B:150:GLY:O	2.36	0.50
2:Y:82:LEU:HD23	2:Y:84:ILE:HD11	1.92	0.50
1:A:43:PHE:O	1:A:151:ARG:NH2	2.33	0.49
1:B:133:ILE:HA	1:B:138:ILE:O	2.12	0.49
1:B:116:TYR:O	1:B:118:VAL:HG13	2.12	0.49
2:Y:44:VAL:HG23	2:Y:48:VAL:HG11	1.95	0.49
1:A:16:LYS:HA	1:A:29:ILE:O	2.13	0.48
1:A:120:LYS:HB3	1:A:127:PHE:CE1	2.48	0.48
1:A:147:LEU:N	1:A:148:PRO:HD2	2.29	0.48
1:B:50:PHE:O	2:Y:97:GLY:N	2.46	0.48
2:X:35:SER:HA	2:X:119:GLN:NE2	2.28	0.48
2:X:74:ASP:OD1	2:X:74:ASP:N	2.44	0.48
2:X:44:VAL:HG11	2:X:115:ALA:HB3	1.95	0.47
2:X:98:GLY:HA2	3:X:1:PO4:O2	2.14	0.47
1:A:139:LEU:HD21	1:A:142:ILE:HD11	1.97	0.47
2:X:37:ARG:NH1	2:X:119:GLN:HG3	2.30	0.47
1:B:130:LEU:C	1:B:130:LEU:HD23	2.40	0.47
1:B:14:ASN:OD1	1:B:31:LEU:HB3	2.15	0.46
1:A:81:HIS:H	1:A:81:HIS:CD2	2.31	0.46
2:Y:66:VAL:HG13	2:Y:108:LEU:HD21	1.97	0.46
1:A:20:VAL:HG21	1:A:84:HIS:HB2	1.97	0.46
2:X:51:ARG:HD2	2:X:102:TYR:CE2	2.51	0.46
1:A:38:TYR:O	1:A:71:SER:HA	2.15	0.46
1:A:181:SER:CB	1:A:182:ASP:HA	2.18	0.45
1:A:181:SER:HA	1:A:183:THR:N	2.20	0.45
1:A:174:PRO:HA	2:Y:91:ASP:O	2.17	0.45
1:A:183:THR:HG22	2:Y:42:HIS:HB3	1.98	0.45
1:B:187:ASP:OD1	1:B:187:ASP:C	2.60	0.45
1:B:32:SER:C	1:B:34:TYR:H	2.25	0.44
1:A:173:SER:HB2	1:A:174:PRO:HD2	1.99	0.44
1:A:100:MET:HA	1:A:100:MET:HE2	1.99	0.44
1:A:141:GLN:NE2	1:A:143:THR:HG22	2.26	0.44
1:A:47:ASP:OD2	1:A:84:HIS:ND1	2.47	0.43
1:A:23:ASP:OD1	1:A:25:GLN:HB2	2.19	0.43
1:B:51:VAL:HG23	2:Y:128:TYR:HE2	1.83	0.43
1:B:34:TYR:CE2	1:B:72:GLN:HG2	2.52	0.43
1:A:52:CYS:C	1:A:54:THR:H	2.27	0.43
2:X:45:PRO:HA	2:X:111:GLU:O	2.18	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:X:122:LEU:O	2:X:125:LEU:HB2	2.19	0.43
2:Y:101:ARG:N	3:Y:2:PO4:O2	2.46	0.43
1:B:47:ASP:O	1:B:87:TRP:CZ2	2.72	0.43
2:X:126:ARG:HA	2:X:133:THR:OG1	2.18	0.43
2:Y:109:GLN:H	2:Y:109:GLN:CD	2.27	0.43
1:B:106:SER:C	1:B:108:PRO:HD3	2.44	0.42
1:A:47:ASP:O	1:A:87:TRP:CD2	2.72	0.42
1:A:174:PRO:HG3	2:Y:90:GLY:HA2	2.00	0.42
2:X:44:VAL:HA	2:X:45:PRO:HD3	1.86	0.42
2:Y:46:LEU:HD22	2:Y:110:ARG:O	2.20	0.42
2:X:52:PRO:HG2	2:X:96:PHE:HE1	1.85	0.42
1:B:90:THR:HA	1:B:91:PRO:HD3	1.87	0.41
1:B:79:ASP:HB2	1:B:84:HIS:CE1	2.55	0.41
1:B:100:MET:HG3	1:B:104:LEU:HD11	2.02	0.41
1:A:140:ARG:HH11	1:A:140:ARG:CB	2.32	0.41
1:B:13:PRO:HD2	1:B:116:TYR:CZ	2.55	0.41
2:X:53:LEU:HA	2:X:54:PRO:HD3	1.86	0.41
2:Y:133:THR:HA	2:Y:134:PRO:HD3	1.86	0.41
1:A:31:LEU:HD21	1:A:133:ILE:HD13	2.03	0.41
2:X:70:ARG:HG2	2:X:71:GLU:HG2	2.01	0.41
1:A:174:PRO:HD3	2:Y:92:TYR:OH	2.21	0.41
1:A:76:ALA:HB2	1:A:105:VAL:HB	2.03	0.40
1:A:106:SER:OG	1:A:108:PRO:HD3	2.21	0.40
1:A:149:VAL:HG21	1:B:163:PHE:CZ	2.56	0.40
1:B:16:LYS:HG3	1:B:29:ILE:O	2.21	0.40
1:B:141:GLN:CD	1:B:143:THR:HG23	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/199 (92%)	166 (91%)	13 (7%)	3 (2%)	8	17
1	B	183/199 (92%)	173 (94%)	8 (4%)	2 (1%)	12	26
2	X	104/110 (94%)	95 (91%)	7 (7%)	2 (2%)	6	13
2	Y	97/110 (88%)	85 (88%)	10 (10%)	2 (2%)	5	11
All	All	566/618 (92%)	519 (92%)	38 (7%)	9 (2%)	8	17

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	X	39	ALA
2	Y	90	GLY
2	Y	98	GLY
1	A	54	THR
1	A	13	PRO
1	A	85	LEU
1	B	182	ASP
2	X	35	SER
1	B	22	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/171 (91%)	132 (85%)	24 (15%)	2	4
1	B	157/171 (92%)	140 (89%)	17 (11%)	5	10
2	X	91/94 (97%)	82 (90%)	9 (10%)	6	13
2	Y	85/94 (90%)	80 (94%)	5 (6%)	16	35
All	All	489/530 (92%)	434 (89%)	55 (11%)	5	9

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	SER
1	A	7	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	18	THR
1	A	30	SER
1	A	35	LYS
1	A	46	LEU
1	A	49	THR
1	A	57	ILE
1	A	62	ARG
1	A	67	LYS
1	A	68	LYS
1	A	70	ASN
1	A	85	LEU
1	A	102	ILE
1	A	122	ASP
1	A	126	SER
1	A	140	ARG
1	A	141	GLN
1	A	143	THR
1	A	147	LEU
1	A	151	ARG
1	A	158	ARG
1	A	166	THR
1	A	178	LYS
1	B	18	THR
1	B	46	LEU
1	B	54	THR
1	B	67	LYS
1	B	83	GLU
1	B	85	LEU
1	B	119	LEU
1	B	122	ASP
1	B	123	GLU
1	B	128	ARG
1	B	141	GLN
1	B	147	LEU
1	B	158	ARG
1	B	178	LYS
1	B	183	THR
1	B	185	LYS
1	B	187	ASP
2	X	37	ARG
2	X	38	ILE
2	X	44	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	X	46	LEU
2	X	48	VAL
2	X	51	ARG
2	X	74	ASP
2	X	121	THR
2	X	123	SER
2	Y	44	VAL
2	Y	82	LEU
2	Y	106	GLN
2	Y	109	GLN
2	Y	117	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	14	ASN
1	A	70	ASN
1	A	72	GLN
1	A	89	ASN
1	A	114	GLN
1	A	141	GLN
1	A	164	GLN
1	B	70	ASN
1	B	101	ASN
1	B	145	ASN
2	X	34	HIS
2	X	119	GLN
2	Y	88	GLN
2	Y	106	GLN
2	Y	109	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	Y	2	-	4,4,4	0.97	0	6,6,6	1.20	1 (16%)
3	PO4	X	1	-	4,4,4	0.99	0	6,6,6	0.41	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	Y	2	PO4	O4-P-O3	2.23	114.85	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Y	2	PO4	1	0
3	X	1	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	184/199 (92%)	-0.38	0 <b>100</b> <b>100</b>	18, 30, 37, 39	0
1	B	185/199 (92%)	-0.43	4 (2%) 62 57	14, 24, 34, 39	0
2	X	106/110 (96%)	-0.44	1 (0%) 81 77	15, 23, 42, 47	0
2	Y	99/110 (90%)	-0.39	1 (1%) 79 75	20, 28, 33, 39	0
All	All	574/618 (92%)	-0.41	6 (1%) 79 75	14, 27, 36, 47	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	X	34	HIS	2.7
1	B	53	PRO	2.3
1	B	50	PHE	2.3
1	B	52	CYS	2.2
1	B	49	THR	2.1
2	Y	109	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	Y	2	5/5	0.74	0.14	67,67,69,70	0
3	PO4	X	1	5/5	0.77	0.11	79,79,80,81	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.