



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2024 – 02:25 PM EDT

PDB ID : 6T36  
Title : Crystal structure of the PTPN3 PDZ domain bound to the HBV core protein C-terminal peptide  
Authors : Genera, M.; Mechaly, A.; Haouz, A.; Caillet-Saguy, C.  
Deposited on : 2019-10-10  
Resolution : 1.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	913	
2	B	212	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 880 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	93	Total	C	N	O	S	0	2	0
			748	467	135	143	3			

- Molecule 2 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	S	0	0	0
			63	33	13	16	1			

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Br	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	51	Total	O	0	1
			52	52		
4	B	16	Total	O	0	0
			16	16		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 3



VAL  
GLN  
MET  
LEU  
ASP  
PRO  
SER

● Molecule 2: Capsid protein

Chain B:  96%

MET GLN THR ASP  
LEU PHE SER  
HIS ALA  
LEU LEU CYS  
CYS LEU  
ILE ILE  
SER LEU  
CYS GLU  
SER SER  
CYS PRO  
THR HIS  
VAL CYS  
GLN SER  
ALA PRO  
SER HIS  
LYS HIS  
LEU THR  
CYS ALA  
LEU LEU  
GLY ARG  
TRP GLN

ASP THR  
ALA  
SER  
LYS  
LEU  
TYR  
ARG  
GLU  
ALA  
LEU  
TRP  
PHE  
HIS  
SER  
PRO  
GLU  
HIS  
CYS  
THR  
CYS  
SER  
PHE  
THR  
GLY  
ARG  
GLU  
THR  
VAL  
LEU  
GLU  
TYR  
ILE  
LEU  
VAL  
SER  
PHE  
GLY  
GLU  
LEU  
TRP  
MET  
TYR  
LYS  
ASN  
LEU  
PHE  
GLY  
ALA  
THR  
TRP  
ALA  
VAL  
SER  
GLY  
VAL  
SER  
ASN  
LEU  
GLU  
SER  
ASP  
PHE  
LEU  
PHE  
PRO  
VAL  
SER  
ILE  
TYR  
VAL  
ASN  
LEU

ASN MET  
GLY  
SER  
LYS  
ILE  
ARG  
GLN  
LEU  
LEU  
TRP  
PHE  
HIS  
SER  
ILE  
SER  
CYS  
LEU  
THR  
PHE  
GLY  
ARG  
GLU  
THR  
ARG  
VAL  
LEU  
GLU  
TYR  
ILE  
LEU  
VAL  
SER  
PHE  
GLY  
VAL  
TRP  
ILE  
ARG  
THR  
PRO  
PRO  
ALA  
TYR  
ARG  
PRO  
GLN  
ASN  
ALA  
PRO  
ILE  
LEU  
SER  
THR  
LEU  
PRO  
GLU  
THR  
THR  
VAL  
VAL  
ARG  
ARG

ARG  
GLY  
ARG  
SER  
PRO  
ARG  
ARG  
ARG  
THR  
PRO  
SER  
PRO  
ARG  
ARG  
ARG  
SER  
GLN  
SER  
PRO  
ARG  
ARG  
ARG  
ARG  
S1  
Q2  
CB

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.58Å 75.58Å 46.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.35 – 1.86 37.96 – 1.86	Depositor EDS
% Data completeness (in resolution range)	98.6 (29.35-1.86) 91.7 (37.96-1.86)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 1.85Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, $R_{free}$	0.195 , 0.239 0.196 , 0.244	Depositor DCC
$R_{free}$ test set	656 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtriage
Anisotropy	0.871	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.049 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	880	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/765	0.57	0/1031
2	B	0.29	0/62	0.44	0/79
All	All	0.34	0/827	0.56	0/1110

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	748	0	759	8	0
2	B	63	0	57	0	0
3	A	1	0	0	0	0
4	A	52	0	0	1	0
4	B	16	0	0	0	0
All	All	880	0	816	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 8 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:563:ASN:HD21	1:A:582:SER:HB3	1.51	0.73
1:A:525:LEU:HD13	1:A:535:LEU:HD11	1.85	0.58
1:A:512:ARG:NH1	4:A:1101:HOH:O	2.30	0.51
1:A:579:ILE:O	1:A:582:SER:OG	2.28	0.51
1:A:582:SER:O	1:A:585:SER:HB3	2.11	0.51

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	93/913 (10%)	91 (98%)	2 (2%)	0	100	100
2	B	6/212 (3%)	6 (100%)	0	0	100	100
All	All	99/1125 (9%)	97 (98%)	2 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/824 (10%)	85 (99%)	1 (1%)	71	62
2	B	8/192 (4%)	8 (100%)	0	100	100
All	All	94/1016 (9%)	93 (99%)	1 (1%)	73	65



All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	505	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	93/913 (10%)	-0.09	3 (3%) 47 45	29, 38, 64, 78	0
2	B	8/212 (3%)	0.18	1 (12%) 3 4	33, 41, 57, 58	0
All	All	101/1125 (8%)	-0.07	4 (3%) 38 36	29, 38, 64, 78	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	586	HIS	5.0
1	A	584	GLU	3.4
1	A	505	ASP	2.9
2	B	2	GLN	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BR	A	1001	1/1	0.99	0.07	50,50,50,50	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.