



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 30, 2026 – 10:16 am BST

PDB ID : 9TJV / pdb_00009tjv
Title : Ternary complex of E. coli leucyl-tRNA synthetase mutant W177A, tRNA(leu) and the leucyl adenylate analog LeuAMS in the aminoacyl transfer state
Authors : Hoffmann, G.; Palencia, A.
Deposited on : 2025-12-08
Resolution : 2.28 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)
RNA backbone	3983	1151 (2.60-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	880	2% 89% 7% .
1	C	880	18% 89% 8% .
2	B	87	% 77% 13% 9% .
2	D	87	80% 7% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 17600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leucine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	852	Total	C	N	O	S	0	0	0
			6774	4303	1149	1280	42			
1	C	851	Total	C	N	O	S	0	0	0
			6760	4295	1146	1277	42			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP C4ZWC9
A	-18	GLY	-	expression tag	UNP C4ZWC9
A	-17	SER	-	expression tag	UNP C4ZWC9
A	-16	SER	-	expression tag	UNP C4ZWC9
A	-15	HIS	-	expression tag	UNP C4ZWC9
A	-14	HIS	-	expression tag	UNP C4ZWC9
A	-13	HIS	-	expression tag	UNP C4ZWC9
A	-12	HIS	-	expression tag	UNP C4ZWC9
A	-11	HIS	-	expression tag	UNP C4ZWC9
A	-10	HIS	-	expression tag	UNP C4ZWC9
A	-9	SER	-	expression tag	UNP C4ZWC9
A	-8	SER	-	expression tag	UNP C4ZWC9
A	-7	GLY	-	expression tag	UNP C4ZWC9
A	-6	LEU	-	expression tag	UNP C4ZWC9
A	-5	VAL	-	expression tag	UNP C4ZWC9
A	-4	PRO	-	expression tag	UNP C4ZWC9
A	-3	ARG	-	expression tag	UNP C4ZWC9
A	-2	GLY	-	expression tag	UNP C4ZWC9
A	-1	SER	-	expression tag	UNP C4ZWC9
A	0	HIS	-	expression tag	UNP C4ZWC9
A	177	ALA	TRP	engineered mutation	UNP C4ZWC9
C	-19	MET	-	initiating methionine	UNP C4ZWC9
C	-18	GLY	-	expression tag	UNP C4ZWC9
C	-17	SER	-	expression tag	UNP C4ZWC9
C	-16	SER	-	expression tag	UNP C4ZWC9

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-15	HIS	-	expression tag	UNP C4ZWC9
C	-14	HIS	-	expression tag	UNP C4ZWC9
C	-13	HIS	-	expression tag	UNP C4ZWC9
C	-12	HIS	-	expression tag	UNP C4ZWC9
C	-11	HIS	-	expression tag	UNP C4ZWC9
C	-10	HIS	-	expression tag	UNP C4ZWC9
C	-9	SER	-	expression tag	UNP C4ZWC9
C	-8	SER	-	expression tag	UNP C4ZWC9
C	-7	GLY	-	expression tag	UNP C4ZWC9
C	-6	LEU	-	expression tag	UNP C4ZWC9
C	-5	VAL	-	expression tag	UNP C4ZWC9
C	-4	PRO	-	expression tag	UNP C4ZWC9
C	-3	ARG	-	expression tag	UNP C4ZWC9
C	-2	GLY	-	expression tag	UNP C4ZWC9
C	-1	SER	-	expression tag	UNP C4ZWC9
C	0	HIS	-	expression tag	UNP C4ZWC9
C	177	ALA	TRP	engineered mutation	UNP C4ZWC9

- Molecule 2 is a RNA chain called tRNA(leu).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	79	Total	C	N	O	P	0	0	0
			1675	743	303	550	79			
2	D	77	Total	C	N	O	P	0	0	0
			1632	724	295	536	77			

- Molecule 3 is 5'-O-(L-leucylsulfamoyl)adenosine (CCD ID: LSS) (formula: C₁₆H₂₅N₇O₇S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 16	N 7	O 7	S 1	0	0
3	C	1	Total 31	C 16	N 7	O 7	S 1	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Mg 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	458	Total O 458 458	0	0
5	B	178	Total O 178 178	0	0
5	C	45	Total O 45 45	0	0
5	D	14	Total O 14 14	0	0

- Molecule 1: Leucine-tRNA ligase






● Molecule 2: tRNA(Leu)



● Molecule 2: tRNA(Leu)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	158.01Å 68.31Å 226.28Å 90.00° 104.44° 90.00°	Depositor
Resolution (Å)	219.13 – 2.28 219.13 – 2.28	Depositor EDS
% Data completeness (in resolution range)	78.3 (219.13-2.28) 78.3 (219.13-2.28)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.194 , 0.222 0.187 , 0.218	Depositor DCC
R_{free} test set	4265 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17600	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LSS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	2/6933 (0.0%)	1.01	13/9410 (0.1%)
1	C	0.57	1/6918 (0.0%)	0.97	15/9390 (0.2%)
2	B	0.61	1/1869 (0.1%)	0.70	0/2908
2	D	0.49	0/1821	0.65	0/2833
All	All	0.62	4/17541 (0.0%)	0.92	28/24541 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	568	MET	SD-CE	-10.07	1.54	1.79
1	A	494	MET	SD-CE	-6.20	1.64	1.79
1	C	568	MET	SD-CE	-5.38	1.66	1.79
2	B	47(D)	C	C1'-N1	5.13	1.55	1.47

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	438	GLU	CA-C-N	8.46	131.96	120.38
1	A	438	GLU	C-N-CA	8.46	131.96	120.38
1	C	438	GLU	CA-C-N	7.28	130.36	120.38
1	C	438	GLU	C-N-CA	7.28	130.36	120.38
1	C	541	PHE	CA-CB-CG	6.69	120.49	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6774	0	6634	38	0
1	C	6760	0	6623	35	0
2	B	1675	0	847	4	0
2	D	1632	0	825	1	0
3	A	31	0	25	1	0
3	C	31	0	25	1	0
4	B	2	0	0	0	0
5	A	458	0	0	1	0
5	B	178	0	0	0	0
5	C	45	0	0	0	0
5	D	14	0	0	0	0
All	All	17600	0	14979	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:610:HIS:HE1	1:C:93:ASN:HB3	1.54	0.72
1:A:610:HIS:CE1	1:C:93:ASN:HB3	2.37	0.59
1:A:54:ARG:HD3	1:A:566:GLN:OE1	2.03	0.59
1:C:588:PRO:HG2	1:C:628:ILE:HD11	1.85	0.58
1:C:54:ARG:HD3	1:C:566:GLN:OE1	2.05	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	848/880 (96%)	835 (98%)	13 (2%)	0	100	100
1	C	847/880 (96%)	831 (98%)	16 (2%)	0	100	100
All	All	1695/1760 (96%)	1666 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	717/740 (97%)	713 (99%)	4 (1%)	78	87
1	C	715/740 (97%)	712 (100%)	3 (0%)	84	91
All	All	1432/1480 (97%)	1425 (100%)	7 (0%)	81	89

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	818	VAL
1	C	231	GLU
1	C	818	VAL
1	C	386	GLU
1	A	744	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	690	ASN
1	C	774	GLN
1	A	805	GLN
1	C	15	GLN
1	C	157	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	75/87 (86%)	7 (9%)	0
2	D	73/87 (83%)	6 (8%)	0
All	All	148/174 (85%)	13 (8%)	0

5 of 13 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	9	G
2	B	13	G
2	B	16	U
2	B	18	G
2	B	19	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	LSS	A	901	-	33,33,33	0.20	0	46,49,49	0.37	0
3	LSS	C	901	-	33,33,33	0.16	0	46,49,49	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LSS	A	901	-	-	2/22/39/39	0/3/3/3
3	LSS	C	901	-	-	2/22/39/39	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	901	LSS	C25-O5-S1-N2
3	C	901	LSS	C25-O5-S1-N2
3	A	901	LSS	C23-C24-C25-O5
3	C	901	LSS	C23-C24-C25-O5

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	LSS	1	0
3	C	901	LSS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	852/880 (96%)	-0.16	14 (1%) 70 72	21, 35, 61, 86	0
1	C	851/880 (96%)	1.21	162 (19%) 3 3	52, 90, 140, 154	0
2	B	79/87 (90%)	-0.71	1 (1%) 75 76	25, 39, 89, 102	0
2	D	77/87 (88%)	0.46	0 100 100	77, 116, 146, 153	0
All	All	1859/1934 (96%)	0.47	177 (9%) 14 15	21, 60, 134, 154	0

The worst 5 of 177 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	608	ALA	6.1
1	C	835	VAL	5.2
1	C	607	ALA	4.9
1	C	737	ALA	4.9
1	C	264	GLY	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	B	101	1/1	0.88	0.17	51,51,51,51	0
4	MG	B	102	1/1	0.94	0.20	45,45,45,45	0
3	LSS	C	901	31/31	0.96	0.08	57,58,59,60	0
3	LSS	A	901	31/31	0.98	0.05	19,23,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LSS C 901:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.