



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2024 – 04:31 PM EDT

PDB ID : 4TQV
Title : Crystal structure of a bacterial ABC transporter involved in the import of the acidic polysaccharide alginate
Authors : Maruyama, Y.; Itoh, T.; Kaneko, A.; Nishitani, Y.; Mikami, B.; Hashimoto, W.; Murata, K.
Deposited on : 2014-06-12
Resolution : 4.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.














Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1055 (5.20-3.80)
Clashscore	141614	1123 (5.20-3.80)
Ramachandran outliers	138981	1069 (5.20-3.80)
Sidechain outliers	138945	1050 (5.20-3.80)
RSRZ outliers	127900	1101 (5.30-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	
1	E	301	
1	I	301	
1	M	301	
2	B	305	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	305	
2	J	305	
2	N	305	
3	C	363	
3	D	363	
3	G	363	
3	H	363	
3	K	363	
3	L	363	
3	O	363	
3	P	363	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 40472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AlgM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	0	0
			2314	1550	368	386	10			
1	E	286	Total	C	N	O	S	0	0	0
			2314	1550	368	386	10			
1	I	286	Total	C	N	O	S	0	0	0
			2314	1550	368	386	10			
1	M	286	Total	C	N	O	S	0	0	0
			2314	1550	368	386	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	expression tag	UNP Q9KWT8
E	24	MET	-	expression tag	UNP Q9KWT8
I	24	MET	-	expression tag	UNP Q9KWT8
M	24	MET	-	expression tag	UNP Q9KWT8

- Molecule 2 is a protein called AlgM2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	284	Total	C	N	O	S	0	0	0
			2257	1506	359	379	13			
2	F	280	Total	C	N	O	S	0	0	0
			2229	1488	354	375	12			
2	J	284	Total	C	N	O	S	0	0	0
			2257	1506	359	379	13			
2	N	284	Total	C	N	O	S	0	0	0
			2257	1506	359	379	13			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	294	LEU	-	expression tag	UNP Q9KWT7
B	295	GLU	-	expression tag	UNP Q9KWT7
B	296	HIS	-	expression tag	UNP Q9KWT7
B	297	HIS	-	expression tag	UNP Q9KWT7
B	298	HIS	-	expression tag	UNP Q9KWT7
B	299	HIS	-	expression tag	UNP Q9KWT7
B	300	HIS	-	expression tag	UNP Q9KWT7
B	301	HIS	-	expression tag	UNP Q9KWT7
B	302	HIS	-	expression tag	UNP Q9KWT7
B	303	HIS	-	expression tag	UNP Q9KWT7
B	304	HIS	-	expression tag	UNP Q9KWT7
B	305	HIS	-	expression tag	UNP Q9KWT7
F	294	LEU	-	expression tag	UNP Q9KWT7
F	295	GLU	-	expression tag	UNP Q9KWT7
F	296	HIS	-	expression tag	UNP Q9KWT7
F	297	HIS	-	expression tag	UNP Q9KWT7
F	298	HIS	-	expression tag	UNP Q9KWT7
F	299	HIS	-	expression tag	UNP Q9KWT7
F	300	HIS	-	expression tag	UNP Q9KWT7
F	301	HIS	-	expression tag	UNP Q9KWT7
F	302	HIS	-	expression tag	UNP Q9KWT7
F	303	HIS	-	expression tag	UNP Q9KWT7
F	304	HIS	-	expression tag	UNP Q9KWT7
F	305	HIS	-	expression tag	UNP Q9KWT7
J	294	LEU	-	expression tag	UNP Q9KWT7
J	295	GLU	-	expression tag	UNP Q9KWT7
J	296	HIS	-	expression tag	UNP Q9KWT7
J	297	HIS	-	expression tag	UNP Q9KWT7
J	298	HIS	-	expression tag	UNP Q9KWT7
J	299	HIS	-	expression tag	UNP Q9KWT7
J	300	HIS	-	expression tag	UNP Q9KWT7
J	301	HIS	-	expression tag	UNP Q9KWT7
J	302	HIS	-	expression tag	UNP Q9KWT7
J	303	HIS	-	expression tag	UNP Q9KWT7
J	304	HIS	-	expression tag	UNP Q9KWT7
J	305	HIS	-	expression tag	UNP Q9KWT7
N	294	LEU	-	expression tag	UNP Q9KWT7
N	295	GLU	-	expression tag	UNP Q9KWT7
N	296	HIS	-	expression tag	UNP Q9KWT7
N	297	HIS	-	expression tag	UNP Q9KWT7
N	298	HIS	-	expression tag	UNP Q9KWT7
N	299	HIS	-	expression tag	UNP Q9KWT7
N	300	HIS	-	expression tag	UNP Q9KWT7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
N	301	HIS	-	expression tag	UNP Q9KWT7
N	302	HIS	-	expression tag	UNP Q9KWT7
N	303	HIS	-	expression tag	UNP Q9KWT7
N	304	HIS	-	expression tag	UNP Q9KWT7
N	305	HIS	-	expression tag	UNP Q9KWT7

- Molecule 3 is a protein called AlgS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	D	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	G	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	H	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	K	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	L	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	O	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			
3	P	363	Total	C	N	O	S	0	0	0
			2777	1745	503	518	11			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	160	GLN	GLU	engineered mutation	UNP Q9KWT9
D	160	GLN	GLU	engineered mutation	UNP Q9KWT9
G	160	GLN	GLU	engineered mutation	UNP Q9KWT9
H	160	GLN	GLU	engineered mutation	UNP Q9KWT9
K	160	GLN	GLU	engineered mutation	UNP Q9KWT9
L	160	GLN	GLU	engineered mutation	UNP Q9KWT9
O	160	GLN	GLU	engineered mutation	UNP Q9KWT9
P	160	GLN	GLU	engineered mutation	UNP Q9KWT9


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AlgM1




• Molecule 1: AlgM1




• Molecule 1: AlgM1






• Molecule 1: AlgM1




• Molecule 2: AlgM2




• Molecule 2: AlgM2







• Molecule 2: AlgM2




• Molecule 2: AlgM2




• Molecule 3: AlgS






• Molecule 3: Algs




• Molecule 3: Algs




• Molecule 3: Algs







• Molecule 3: AlgS




• Molecule 3: AlgS



• Molecule 3: AlgS



• Molecule 3: AlgS



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	115.25Å 151.19Å 162.41Å 68.66° 81.76° 90.10°	Depositor
Resolution (Å)	29.82 – 4.50 29.82 – 4.50	Depositor EDS
% Data completeness (in resolution range)	95.6 (29.82-4.50) 80.3 (29.82-4.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 4.42Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.279 , 0.320 0.281 , 0.320	Depositor DCC
R_{free} test set	2000 reflections (3.50%)	wwPDB-VP
Wilson B-factor (Å ²)	169.5	Xtriage
Anisotropy	0.225	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.20 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	40472	wwPDB-VP
Average B, all atoms (Å ²)	159.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/2377	0.48	1/3241 (0.0%)
1	E	0.25	0/2377	0.52	1/3241 (0.0%)
1	I	0.24	0/2377	0.47	1/3241 (0.0%)
1	M	0.24	0/2377	0.49	2/3241 (0.1%)
2	B	0.26	0/2316	0.50	1/3149 (0.0%)
2	F	0.26	0/2288	0.52	1/3113 (0.0%)
2	J	0.26	0/2316	0.50	1/3149 (0.0%)
2	N	0.28	0/2316	0.53	2/3149 (0.1%)
3	C	0.25	0/2822	0.56	0/3826
3	D	0.26	0/2822	0.60	0/3826
3	G	0.26	0/2822	0.59	0/3826
3	H	0.25	0/2822	0.61	2/3826 (0.1%)
3	K	0.26	0/2822	0.57	0/3826
3	L	0.25	0/2822	0.60	0/3826
3	O	0.26	0/2822	0.56	0/3826
3	P	0.26	0/2822	0.62	2/3826 (0.1%)
All	All	0.26	0/41320	0.55	14/56132 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
1	I	0	1
1	M	0	1
3	C	0	1
3	K	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	298	ARG	NE-CZ-NH1	-7.79	116.41	120.30
2	N	282	PHE	CB-CA-C	-6.60	97.20	110.40
2	J	282	PHE	CB-CA-C	-5.93	98.55	110.40
3	H	298	ARG	NE-CZ-NH1	-5.84	117.38	120.30
3	P	298	ARG	CA-CB-CG	5.56	125.64	113.40

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	77	TRP	Peptide
3	C	21	HIS	Mainchain
1	E	77	TRP	Peptide
1	I	77	TRP	Peptide
3	K	21	HIS	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2314	0	2402	81	1
1	E	2314	0	2402	82	0
1	I	2314	0	2402	72	1
1	M	2314	0	2402	79	0
2	B	2257	0	2320	63	0
2	F	2229	0	2286	60	1
2	J	2257	0	2320	62	0
2	N	2257	0	2320	71	1
3	C	2777	0	2854	76	0
3	D	2777	0	2854	77	0
3	G	2777	0	2854	90	0
3	H	2777	0	2854	90	0
3	K	2777	0	2854	71	0
3	L	2777	0	2854	84	0
3	O	2777	0	2854	70	0
3	P	2777	0	2854	86	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	40472	0	41686	1126	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:34:ASP:OD1	2:N:106:ARG:NH1	1.72	1.23
1:E:34:ASP:OD1	2:F:106:ARG:NH1	2.02	0.92
2:N:102:LEU:HD12	2:N:161:ARG:HE	1.40	0.87
3:L:195:GLN:HG3	3:L:235:ILE:HA	1.59	0.84
2:N:76:SER:HB3	2:N:228:LEU:H	1.42	0.84

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:289:ARG:NH1	2:N:241:VAL:O[1_654]	2.08	0.12
1:A:289:ARG:NH1	2:F:241:VAL:O[1_654]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	282/301 (94%)	257 (91%)	24 (8%)	1 (0%)	34 72
1	E	282/301 (94%)	252 (89%)	26 (9%)	4 (1%)	11 47
1	I	282/301 (94%)	257 (91%)	24 (8%)	1 (0%)	34 72
1	M	282/301 (94%)	257 (91%)	24 (8%)	1 (0%)	34 72
2	B	282/305 (92%)	249 (88%)	31 (11%)	2 (1%)	22 62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	278/305 (91%)	243 (87%)	32 (12%)	3 (1%)	14	52
2	J	282/305 (92%)	249 (88%)	31 (11%)	2 (1%)	22	62
2	N	282/305 (92%)	249 (88%)	31 (11%)	2 (1%)	22	62
3	C	361/363 (99%)	323 (90%)	32 (9%)	6 (2%)	9	43
3	D	361/363 (99%)	320 (89%)	31 (9%)	10 (3%)	5	33
3	G	361/363 (99%)	321 (89%)	35 (10%)	5 (1%)	11	47
3	H	361/363 (99%)	320 (89%)	31 (9%)	10 (3%)	5	33
3	K	361/363 (99%)	323 (90%)	32 (9%)	6 (2%)	9	43
3	L	361/363 (99%)	320 (89%)	31 (9%)	10 (3%)	5	33
3	O	361/363 (99%)	323 (90%)	32 (9%)	6 (2%)	9	43
3	P	361/363 (99%)	320 (89%)	31 (9%)	10 (3%)	5	33
All	All	5140/5328 (96%)	4583 (89%)	478 (9%)	79 (2%)	10	46

5 of 79 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	78	ILE
3	C	18	THR
3	C	322	GLY
3	C	336	VAL
3	C	359	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/260 (96%)	218 (87%)	32 (13%)	4	21
1	E	250/260 (96%)	217 (87%)	33 (13%)	4	20
1	I	250/260 (96%)	218 (87%)	32 (13%)	4	21
1	M	250/260 (96%)	219 (88%)	31 (12%)	4	22
2	B	240/258 (93%)	220 (92%)	20 (8%)	11	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	237/258 (92%)	215 (91%)	22 (9%)	9	30
2	J	240/258 (93%)	220 (92%)	20 (8%)	11	36
2	N	240/258 (93%)	221 (92%)	19 (8%)	12	38
3	C	307/307 (100%)	264 (86%)	43 (14%)	3	19
3	D	307/307 (100%)	259 (84%)	48 (16%)	2	16
3	G	307/307 (100%)	263 (86%)	44 (14%)	3	18
3	H	307/307 (100%)	259 (84%)	48 (16%)	2	16
3	K	307/307 (100%)	264 (86%)	43 (14%)	3	19
3	L	307/307 (100%)	259 (84%)	48 (16%)	2	16
3	O	307/307 (100%)	264 (86%)	43 (14%)	3	19
3	P	307/307 (100%)	259 (84%)	48 (16%)	2	16
All	All	4413/4528 (98%)	3839 (87%)	574 (13%)	4	20

5 of 574 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	320	LYS
3	P	359	GLN
2	N	224	GLU
1	M	310	PHE
3	O	321	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
3	K	357	GLN
1	M	220	GLN
3	L	160	GLN
3	L	345	HIS
3	O	160	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	286/301 (95%)	-0.41	1 (0%) 94 90	45, 123, 232, 396	0
1	E	286/301 (95%)	-0.39	6 (2%) 63 54	72, 144, 245, 389	0
1	I	286/301 (95%)	-0.39	1 (0%) 94 90	63, 110, 236, 381	0
1	M	286/301 (95%)	-0.35	4 (1%) 75 66	63, 137, 231, 308	0
2	B	284/305 (93%)	-0.29	6 (2%) 63 54	40, 135, 260, 432	0
2	F	280/305 (91%)	-0.35	5 (1%) 68 59	74, 138, 252, 482	0
2	J	284/305 (93%)	-0.37	7 (2%) 57 48	64, 129, 253, 401	0
2	N	284/305 (93%)	-0.43	5 (1%) 68 59	50, 112, 244, 387	0
3	C	363/363 (100%)	-0.14	10 (2%) 53 43	44, 179, 275, 344	0
3	D	363/363 (100%)	0.11	25 (6%) 16 13	72, 187, 302, 373	0
3	G	363/363 (100%)	0.02	12 (3%) 46 37	70, 160, 285, 349	0
3	H	363/363 (100%)	-0.01	14 (3%) 39 31	46, 146, 267, 367	0
3	K	363/363 (100%)	-0.19	7 (1%) 66 58	47, 156, 263, 338	0
3	L	363/363 (100%)	-0.06	17 (4%) 31 26	69, 154, 276, 340	0
3	O	363/363 (100%)	0.12	24 (6%) 18 15	65, 198, 291, 376	0
3	P	363/363 (100%)	-0.04	12 (3%) 46 37	57, 162, 279, 355	0
All	All	5180/5328 (97%)	-0.18	156 (3%) 50 39	40, 151, 275, 482	0

The worst 5 of 156 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	P	1	MET	12.2
3	D	1	MET	8.8
3	O	1	MET	7.9
3	O	2	VAL	7.1
3	H	296	SER	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.