



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 12, 2024 – 06:13 PM EST

PDB ID : 3U79  
Title : AL-103 Y32F Y96F  
Authors : DiCostanzo, A.C.; Thompson, J.R.; Ramirez-Alvarado, M.  
Deposited on : 2011-10-13  
Resolution : 1.62 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6077 (1.64-1.60)
Clashscore	180529	6617 (1.64-1.60)
Ramachandran outliers	177936	6498 (1.64-1.60)
Sidechain outliers	177891	6497 (1.64-1.60)
RSRZ outliers	164620	6075 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	111	<div> <div>2%</div> <div>81% 19%</div> </div>
1	B	111	<div> <div>2%</div> <div>86% 14%</div> </div>
1	C	111	<div> <div>90% 9%</div> </div>
1	D	111	<div> <div>91% 7%</div> </div>
1	E	111	<div> <div>2%</div> <div>86% 14%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	111	<div><div></div><div>90%</div><div>10%</div></div>
1	G	111	<div><div>4%</div><div></div><div>89%</div><div>10%</div><div></div></div>
1	H	111	<div><div>%</div><div></div><div>88%</div><div>12%</div><div></div></div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15328 atoms, of which 6946 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	111	Total	C	H	N	O	S	0	7	0
			1774	566	881	147	176	4			
1	B	111	Total	C	H	N	O	S	0	5	0
			1761	561	875	146	176	3			
1	C	111	Total	C	H	N	O	S	0	5	0
			1767	562	878	148	176	3			
1	D	110	Total	C	H	N	O	S	0	7	0
			1735	555	861	138	177	4			
1	E	110	Total	C	H	N	O	S	0	5	0
			1732	554	858	141	176	3			
1	F	111	Total	C	H	N	O	S	0	8	0
			1771	565	877	144	182	3			
1	G	110	Total	C	H	N	O	S	0	5	0
			1709	548	845	137	176	3			
1	H	111	Total	C	H	N	O	S	0	2	0
			1690	546	831	137	173	3			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Zn	0	0
			4	4		
2	B	2	Total	Zn	0	0
			2	2		
2	C	4	Total	Zn	0	0
			4	4		
2	D	1	Total	Zn	0	0
			1	1		
2	E	1	Total	Zn	0	0
			1	1		
2	F	1	Total	Zn	0	0
			1	1		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Zn	0	0
			1	1		
2	H	2	Total	Zn	0	0
			2	2		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			7	2	3	2		
3	A	1	Total	C	H	O	0	0
			7	2	3	2		
3	B	1	Total	C	H	O	0	0
			7	2	3	2		
3	C	1	Total	C	H	O	0	0
			7	2	3	2		
3	D	1	Total	C	H	O	0	0
			7	2	3	2		
3	D	1	Total	C	H	O	0	0
			7	2	3	2		
3	E	1	Total	C	H	O	0	0
			7	2	3	2		
3	F	1	Total	C	H	O	0	0
			7	2	3	2		
3	F	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	E	1	Total	C	H	N	O	S	
			25	6	13	1	4	1	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	161	Total	O		
			166	166	0	8
5	B	168	Total	O		
			173	173	0	8
5	C	178	Total	O		
			186	186	0	12
5	D	148	Total	O		
			152	152	0	5
5	E	123	Total	O		
			128	128	0	8
5	F	181	Total	O		
			187	187	0	14
5	G	127	Total	O		
			131	131	0	5
5	H	155	Total	O		
			162	162	0	7



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain



- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain




- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain



- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain




- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain








- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain



- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain



- Molecule 1: Amyloidogenic immunoglobulin light chain protein AL-103 Y32F Y96F, variable domain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.02Å 128.02Å 98.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.06 – 1.62 64.06 – 1.62	Depositor EDS
% Data completeness (in resolution range)	95.5 (64.06-1.62) 96.3 (64.06-1.62)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.62Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.169 , 0.223 0.193 , 0.230	Depositor DCC
$R_{free}$ test set	557 reflections (0.50%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.4	Xtriage
Anisotropy	0.295	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15328	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, ZN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/936	0.44	0/1272
1	B	0.24	0/922	0.43	0/1251
1	C	0.23	0/925	0.42	0/1255
1	D	0.25	0/919	0.43	0/1249
1	E	0.23	0/910	0.41	0/1237
1	F	0.24	0/941	0.42	0/1279
1	G	0.23	0/900	0.43	0/1225
1	H	0.22	0/887	0.42	0/1208
All	All	0.24	0/7340	0.42	0/9976

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	893	881	881	16	0
1	B	886	875	876	7	0
1	C	889	878	878	7	0
1	D	874	861	852	4	0
1	E	874	858	858	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	894	877	865	9	0
1	G	864	845	846	8	0
1	H	859	831	831	8	0
2	A	4	0	0	0	0
2	B	2	0	0	0	0
2	C	4	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	2	0	0	0	0
3	A	8	6	6	0	0
3	B	4	3	3	0	0
3	C	4	3	3	0	0
3	D	8	6	6	1	0
3	E	4	3	3	0	0
3	F	8	6	6	0	0
4	E	12	13	13	2	0
5	A	166	0	0	2	0
5	B	173	0	0	0	0
5	C	186	0	0	1	0
5	D	152	0	0	0	0
5	E	128	0	0	0	0
5	F	187	0	0	0	0
5	G	131	0	0	2	0
5	H	162	0	0	1	0
All	All	8382	6946	6927	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:37:GLN:HB2	1:C:47:LEU:HD11	1.74	0.67
1:G:7:SER:OG	5:G:985:HOH:O	2.15	0.63
1:A:6:GLN:HG2	1:A:23[B]:CYS:SG	2.43	0.59
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.85	0.58
1:C:34:ILE:HG23	1:C:46:LEU:CD1	2.34	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/111 (104%)	111 (96%)	5 (4%)	0	100	100
1	B	114/111 (103%)	109 (96%)	4 (4%)	1 (1%)	14	3
1	C	114/111 (103%)	108 (95%)	6 (5%)	0	100	100
1	D	114/111 (103%)	108 (95%)	6 (5%)	0	100	100
1	E	112/111 (101%)	106 (95%)	6 (5%)	0	100	100
1	F	117/111 (105%)	112 (96%)	5 (4%)	0	100	100
1	G	113/111 (102%)	108 (96%)	5 (4%)	0	100	100
1	H	111/111 (100%)	104 (94%)	7 (6%)	0	100	100
All	All	911/888 (103%)	866 (95%)	44 (5%)	1 (0%)	48	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	93	ASN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/98 (107%)	105 (100%)	0	100	100
1	B	103/98 (105%)	103 (100%)	0	100	100
1	C	103/98 (105%)	102 (99%)	1 (1%)	73	57
1	D	104/98 (106%)	101 (97%)	3 (3%)	37	13

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	102/98 (104%)	100 (98%)	2 (2%)	50	25
1	F	106/98 (108%)	104 (98%)	2 (2%)	52	28
1	G	102/98 (104%)	102 (100%)	0	100	100
1	H	99/98 (101%)	99 (100%)	0	100	100
All	All	824/784 (105%)	816 (99%)	8 (1%)	84	57

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	107[B]	ILE
1	F	107[A]	ILE
1	E	53[A]	ASN
1	D	98	THR
1	E	53[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	53	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 26 ligands modelled in this entry, 16 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	E	111	2	3,3,3	0.81	0	3,3,3	1.36	0
3	ACT	A	113	2	3,3,3	0.82	0	3,3,3	1.36	0
3	ACT	F	113	2	3,3,3	0.81	0	3,3,3	1.39	0
3	ACT	D	111	2	3,3,3	0.82	0	3,3,3	1.33	0
3	ACT	B	112	2	3,3,3	0.82	0	3,3,3	1.34	0
3	ACT	D	112	2	3,3,3	0.81	0	3,3,3	1.32	0
3	ACT	F	112	2	3,3,3	0.83	0	3,3,3	1.35	0
4	MES	E	112	-	12,12,12	0.80	0	15,16,16	2.33	6 (40%)
3	ACT	C	114	2	3,3,3	0.81	0	3,3,3	1.37	0
3	ACT	A	112	2	3,3,3	0.82	0	3,3,3	1.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	E	112	-	-	1/6/14/14	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	112	MES	O1S-S-C8	5.21	114.60	106.73
4	E	112	MES	O3S-S-C8	4.22	114.25	106.00
4	E	112	MES	O2S-S-C8	3.48	111.98	106.73
4	E	112	MES	O2S-S-O1S	-2.73	104.96	113.82
4	E	112	MES	O3S-S-O2S	-2.68	104.70	111.40

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	112	MES	C8-C7-N4-C5

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	111	ACT	1	0
4	E	112	MES	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	111/111 (100%)	-0.36	1 (0%) 81 83	12, 22, 37, 53	7 (6%)
1	B	111/111 (100%)	-0.37	1 (0%) 81 83	11, 22, 37, 47	5 (4%)
1	C	111/111 (100%)	0.11	0 100 100	14, 25, 37, 48	5 (4%)
1	D	110/111 (99%)	-0.26	0 100 100	14, 26, 44, 64	6 (5%)
1	E	110/111 (99%)	0.58	2 (1%) 67 70	17, 31, 44, 53	5 (4%)
1	F	111/111 (100%)	0.11	0 100 100	11, 24, 43, 52	7 (6%)
1	G	110/111 (99%)	0.50	4 (3%) 46 48	14, 30, 49, 64	5 (4%)
1	H	111/111 (100%)	-0.23	1 (0%) 81 83	12, 26, 43, 57	2 (1%)
All	All	885/888 (99%)	0.01	9 (1%) 79 82	11, 26, 45, 64	42 (4%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	109	ARG	4.1
1	G	107	ILE	2.7
1	G	-1	SER	2.6
1	E	3[A]	GLN	2.5
1	A	109	ARG	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MES	E	112	12/12	0.54	0.21	36,52,61,61	0
2	ZN	C	113	1/1	0.60	0.19	48,48,48,48	1
3	ACT	A	113	4/4	0.76	0.13	36,42,43,47	0
2	ZN	A	111	1/1	0.77	0.14	42,42,42,42	1
3	ACT	F	113	4/4	0.78	0.20	31,33,40,40	0
3	ACT	A	112	4/4	0.84	0.16	34,36,43,43	0
3	ACT	E	111	4/4	0.85	0.12	37,40,44,44	0
3	ACT	C	114	4/4	0.86	0.11	36,38,45,45	0
2	ZN	H	111	1/1	0.89	0.12	26,26,26,26	1
3	ACT	F	112	4/4	0.89	0.09	34,35,42,42	0
3	ACT	D	111	4/4	0.91	0.11	23,30,37,37	0
2	ZN	C	112	1/1	0.93	0.09	30,30,30,30	0
2	ZN	B	111	1/1	0.94	0.07	50,50,50,50	1
3	ACT	B	112	4/4	0.94	0.10	27,33,44,44	0
2	ZN	A	114	1/1	0.95	0.13	30,30,30,30	1
2	ZN	C	111	1/1	0.96	0.09	30,30,30,30	0
3	ACT	D	112	4/4	0.96	0.06	23,25,30,30	0
2	ZN	F	110	1/1	0.97	0.08	33,33,33,33	0
2	ZN	A	110	1/1	0.97	0.11	33,33,33,33	1
2	ZN	G	111	1/1	0.98	0.06	33,33,33,33	0
2	ZN	E	110	1/1	0.98	0.07	38,38,38,38	0
2	ZN	D	110	1/1	0.99	0.05	23,23,23,23	0
2	ZN	C	110	1/1	0.99	0.02	20,20,20,20	0
2	ZN	A	115	1/1	0.99	0.03	23,23,23,23	0
2	ZN	H	110	1/1	1.00	0.01	17,17,17,17	0
2	ZN	B	110	1/1	1.00	0.04	27,27,27,27	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.