



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 2, 2024 – 02:13 PM EST

PDB ID : 2VDP
Title : Integrin AlphaIIbBeta3 Headpiece Bound to Fibrinogen Gamma chain peptide,LGGAKQAGDV
Authors : Springer, T.A.; Zhu, J.; Xiao, T.
Deposited on : 2007-10-10
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	452	 93% 7%
2	B	461	 5% 91% 8%
3	C	10	 40% 80% 20%
4	H	221	 2% 90% 9%
5	L	214	 94% 6%
6	D	5	 60% 40%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	E	7	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	NAG	B	3099	X	-	-	-
6	MAN	D	3	X	-	-	-
7	MAN	E	3	X	-	-	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 11208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTEGRIN ALPHA-IIB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	452	Total	C	N	O	S	0	8	0
			3514	2232	611	663	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	282	GLY	ALA	conflict	UNP P08514

- Molecule 2 is a protein called INTEGRIN BETA-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	455	Total	C	N	O	S	0	5	0
			3554	2217	606	697	34			

- Molecule 3 is a protein called FIBRINOGEN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	8	Total	C	N	O	0	0	0
			52	30	10	12			


- Molecule 4 is a protein called MONOCLONAL ANTIBODY 10E5 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	219	Total	C	N	O	S	0	2	0
			1672	1060	269	337	6			

- Molecule 5 is a protein called MONOCLONAL ANTIBODY 10E5 LIGHT CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	L	214	Total	C	N	O	S	0	4	0
			1654	1030	269	346	9			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	D	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	E	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	4	Total	Ca	0	0
			4	4		
9	B	2	Total	Ca	0	0
			2	2		

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			14	8	1	5		
10	A	1	Total	C	N	O	0	0
			14	8	1	5		
10	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 11 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	1	Total 1	Mg 1	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	228	Total 228	O 228	0	0
12	B	102	Total 102	O 102	0	0
12	C	7	Total 7	O 7	0	0
12	H	106	Total 106	O 106	0	0
12	L	114	Total 114	O 114	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: INTEGRIN ALPHA-IIB




- Molecule 2: INTEGRIN BETA-3



- Molecule 3: FIBRINOGEN



- Molecule 4: MONOCLONAL ANTIBODY 10E5 HEAVY CHAIN



- Molecule 5: MONOCLONAL ANTIBODY 10E5 LIGHT CHAIN





- Molecule 6: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] α -D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] α -D-mannopyranose-(1-6)-[α -D-mannopyranose-(1-3)] α -D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	148.43Å 148.43Å 177.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.23 – 2.80 46.23 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.3 (46.23-2.80) 93.6 (46.23-2.80)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.89 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.156 , 0.204 0.162 , 0.210	Depositor DCC
R_{free} test set	2660 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	53.8	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 63.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11208	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA, MAN, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3623	0.53	0/4935
2	B	0.33	0/3626	0.49	0/4913
3	C	0.52	0/51	0.56	0/65
4	H	0.32	0/1721	0.50	0/2357
5	L	0.33	0/1702	0.52	0/2308
All	All	0.33	0/10723	0.51	0/14578

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3514	0	3356	20	0
2	B	3554	0	3493	18	0
3	C	52	0	49	0	0
4	H	1672	0	1629	16	0
5	L	1654	0	1573	7	0
6	D	61	0	52	0	0
7	E	83	0	70	0	0
8	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	6	0	7	0	0
9	A	4	0	0	0	0
9	B	2	0	0	0	0
10	A	28	0	26	0	0
10	B	14	0	13	0	0
11	B	1	0	0	0	0
12	A	228	0	0	1	0
12	B	102	0	0	0	0
12	C	7	0	0	0	0
12	H	106	0	0	1	0
12	L	114	0	0	0	0
All	All	11208	0	10276	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 58 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:LEU:HD23	1:A:276[A]:ARG:HA	1.65	0.78
1:A:270:LEU:HD23	1:A:276[B]:ARG:HA	1.66	0.77
5:L:122:SER:O	5:L:126:THR:HG23	2.01	0.61
1:A:235:TRP:HZ2	1:A:270:LEU:HD11	1.65	0.60
1:A:196:ALA:HA	1:A:221:LEU:HD12	1.84	0.59

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	456/452 (101%)	443 (97%)	12 (3%)	1 (0%)	44 73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	456/461 (99%)	427 (94%)	27 (6%)	2 (0%)	30	61
3	C	6/10 (60%)	5 (83%)	1 (17%)	0	100	100
4	H	217/221 (98%)	210 (97%)	7 (3%)	0	100	100
5	L	216/214 (101%)	211 (98%)	5 (2%)	0	100	100
All	All	1351/1358 (100%)	1296 (96%)	52 (4%)	3 (0%)	44	73

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	9	GLY
1	A	123	GLU
2	B	80	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/360 (102%)	358 (98%)	8 (2%)	47	79
2	B	410/409 (100%)	408 (100%)	2 (0%)	86	95
3	C	4/5 (80%)	4 (100%)	0	100	100
4	H	191/190 (100%)	189 (99%)	2 (1%)	73	91
5	L	192/188 (102%)	191 (100%)	1 (0%)	86	95
All	All	1163/1152 (101%)	1150 (99%)	13 (1%)	70	90

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	413	THR
2	B	215	ASN
5	L	160	LEU
4	H	133	VAL
4	H	183	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	316	ASN
2	B	428	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	D	1	2,6	14,14,15	0.62	0	17,19,21	0.78	0
6	NAG	D	2	6	14,14,15	0.71	0	17,19,21	0.93	1 (5%)
6	MAN	D	3	6	11,11,12	0.58	0	15,15,17	0.65	0
6	MAN	D	4	6	11,11,12	0.55	0	15,15,17	0.60	0
6	MAN	D	5	6	11,11,12	0.59	0	15,15,17	0.85	1 (6%)
7	NAG	E	1	2,7	14,14,15	0.64	0	17,19,21	0.83	0
7	NAG	E	2	7	14,14,15	0.48	0	17,19,21	1.06	1 (5%)
7	MAN	E	3	7	11,11,12	0.54	0	15,15,17	0.92	0
7	MAN	E	4	7	11,11,12	0.58	0	15,15,17	1.14	1 (6%)
7	MAN	E	5	7	11,11,12	0.59	0	15,15,17	1.05	1 (6%)
7	MAN	E	6	7	11,11,12	0.55	0	15,15,17	0.98	1 (6%)
7	MAN	E	7	7	11,11,12	0.58	0	15,15,17	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	D	1	2,6	-	0/6/23/26	0/1/1/1
6	NAG	D	2	6	-	0/6/23/26	0/1/1/1
6	MAN	D	3	6	1/1/4/5	2/2/19/22	0/1/1/1
6	MAN	D	4	6	-	2/2/19/22	0/1/1/1
6	MAN	D	5	6	-	2/2/19/22	0/1/1/1
7	NAG	E	1	2,7	-	0/6/23/26	0/1/1/1
7	NAG	E	2	7	-	2/6/23/26	0/1/1/1
7	MAN	E	3	7	1/1/4/5	1/2/19/22	0/1/1/1
7	MAN	E	4	7	-	0/2/19/22	0/1/1/1
7	MAN	E	5	7	-	0/2/19/22	0/1/1/1
7	MAN	E	6	7	-	0/2/19/22	0/1/1/1
7	MAN	E	7	7	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	4	MAN	C1-O5-C5	3.73	117.18	112.19
7	E	2	NAG	C1-O5-C5	2.98	116.18	112.19
7	E	6	MAN	C1-O5-C5	2.54	115.58	112.19
7	E	5	MAN	C1-O5-C5	2.23	115.18	112.19
6	D	2	NAG	C4-C3-C2	2.22	114.28	111.02

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	D	3	MAN	C1
7	E	3	MAN	C1


5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	D	4	MAN	O5-C5-C6-O6
6	D	4	MAN	C4-C5-C6-O6
6	D	3	MAN	O5-C5-C6-O6
7	E	2	NAG	O5-C5-C6-O6
6	D	5	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GOL	B	1462	9	5,5,5	0.41	0	5,5,5	0.20	0
10	NAG	A	3015	1	14,14,15	0.52	0	17,19,21	0.77	0
10	NAG	B	3099	2	14,14,15	0.73	0	17,19,21	1.24	2 (11%)
8	GOL	A	1453	-	5,5,5	0.45	0	5,5,5	0.24	0
10	NAG	A	3249	1	14,14,15	0.56	0	17,19,21	1.14	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	B	1462	9	-	2/4/4/4	-
10	NAG	A	3015	1	-	2/6/23/26	0/1/1/1
10	NAG	B	3099	2	1/1/5/7	2/6/23/26	0/1/1/1
8	GOL	A	1453	-	-	2/4/4/4	-
10	NAG	A	3249	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	3249	NAG	C1-O5-C5	3.76	117.22	112.19
10	B	3099	NAG	C4-C3-C2	2.91	115.28	111.02
10	B	3099	NAG	O5-C1-C2	2.35	114.92	111.29

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
10	B	3099	NAG	C1

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	1462	GOL	C1-C2-C3-O3
10	B	3099	NAG	O5-C5-C6-O6
10	B	3099	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
10	A	3015	NAG	O5-C5-C6-O6
8	B	1462	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Warning: The R factor obtained from EDS is 0.2176, which does not match the depositor's R factor of 0.156. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/452 (100%)	-0.21	3 (0%) 84 79	22, 51, 70, 85	8 (1%)
2	B	455/461 (98%)	0.39	25 (5%) 32 25	23, 63, 107, 121	8 (1%)
3	C	8/10 (80%)	2.25	4 (50%) 0 1	49, 79, 96, 102	0
4	H	219/221 (99%)	-0.07	4 (1%) 67 60	30, 52, 79, 110	3 (1%)
5	L	214/214 (100%)	-0.29	1 (0%) 87 83	20, 46, 64, 89	3 (1%)
All	All	1348/1358 (99%)	0.02	37 (2%) 56 47	20, 53, 95, 121	22 (1%)

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337[A]	PRO	6.5
3	C	405	ALA	5.7
2	B	72	LYS	4.5
2	B	458	GLY	4.1
2	B	441	ALA	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
10	NAG	A	3249	14/15	0.35	0.20	92,95,96,96	14
10	NAG	B	3099	14/15	0.55	0.18	100,105,110,112	0
10	NAG	A	3015	14/15	0.67	0.21	93,95,96,96	14
8	GOL	B	1462	6/6	0.93	0.15	68,71,72,72	0
8	GOL	A	1453	6/6	0.93	0.13	69,70,73,75	0
9	CA	A	2007	1/1	0.99	0.11	63,63,63,63	0
9	CA	B	2002	1/1	0.99	0.13	49,49,49,49	0
9	CA	B	2003	1/1	0.99	0.15	39,39,39,39	0
9	CA	A	2004	1/1	0.99	0.10	62,62,62,62	0
9	CA	A	2005	1/1	0.99	0.14	57,57,57,57	0
9	CA	A	2006	1/1	0.99	0.09	66,66,66,66	0
11	MG	B	2001	1/1	0.99	0.12	33,33,33,33	0

6.5 Other polymers

There are no such residues in this entry.