



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 12, 2024 – 09:04 AM EDT

PDB ID : 5W3V
Title : Crystal Structure of macaque APOBEC3H in complex with RNA
Authors : Bohn, J.A.; Thummar, K.; York, A.; Raymond, A.; Brown, W.C.; Bieniasz, P.D.; Hatzioannou, T.; Smith, J.L.
Deposited on : 2017-06-08
Resolution : 2.24 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3139 (2.26-2.22)
Clashscore	180529	3381 (2.26-2.22)
Ramachandran outliers	177936	3334 (2.26-2.22)
Sidechain outliers	177891	3335 (2.26-2.22)
RSRZ outliers	164620	3138 (2.26-2.22)
RNA backbone	3690	1058 (2.58-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	<div> <div>2%</div> <div>78% 8% 13%</div> </div>
2	B	215	<div> <div>74% 11% 15%</div> </div>
2	C	215	<div> <div>2%</div> <div>74% 12% 14%</div> </div>
2	D	215	<div> <div>73% 12% 15%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	10	 70%30%
3	H	10	 60%30%10%
4	F	10	 60%30%10%
4	G	10	 70%30%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7204 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Apobec3H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	186	Total	C	N	O	S	0	2	0
			1537	976	274	276	11			

- Molecule 2 is a protein called Apobec3H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	182	Total	C	N	O	S	0	3	0
			1525	968	274	274	9			
2	C	184	Total	C	N	O	S	0	1	0
			1522	969	271	273	9			
2	D	183	Total	C	N	O	S	0	4	0
			1536	973	275	277	11			

- Molecule 3 is a RNA chain called RNA (5'-R(P*AP*AP*CP*CP*CP*GP*GP*GP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	10	Total	C	N	O	P	0	0	0
			209	92	39	68	10			
3	H	10	Total	C	N	O	P	0	0	0
			209	92	39	68	10			

- Molecule 4 is a RNA chain called RNA (5'-R(P*AP*AP*CP*CP*CP*CP*GP*GP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	10	Total	C	N	O	P	0	0	0
			206	91	37	68	10			
4	G	10	Total	C	N	O	P	0	0	0
			206	91	37	68	10			




- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Zn 1	0	0
5	B	1	Total 1	Zn 1	0	0
5	C	1	Total 1	Zn 1	0	0
5	D	1	Total 1	Zn 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	66	Total 66	O 66	0	0
6	B	30	Total 30	O 30	0	0
6	C	33	Total 33	O 33	0	0
6	D	76	Total 76	O 76	0	0
6	E	14	Total 14	O 14	0	0
6	F	9	Total 9	O 9	0	0
6	G	12	Total 12	O 12	0	0
6	H	10	Total 10	O 10	0	0

- Molecule 1: Apobec3H



- Molecule 3: RNA (5'-R(P*AP*AP*CP*CP*CP*GP*GP*GP*GP*A)-3')

Chain E:  70% 30%



- Molecule 3: RNA (5'-R(P*AP*AP*CP*CP*CP*GP*GP*GP*GP*A)-3')

Chain H:  60% 30% 10%



- Molecule 4: RNA (5'-R(P*AP*AP*CP*CP*CP*CP*GP*GP*GP*C)-3')

Chain F:  60% 30% 10%



- Molecule 4: RNA (5'-R(P*AP*AP*CP*CP*CP*CP*GP*GP*GP*C)-3')

Chain G:  70% 30%



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	87.96Å 89.31Å 134.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.66 – 2.24 44.66 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.66-2.24) 91.3 (44.66-2.24)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.52 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.182 , 0.222 0.188 , 0.228	Depositor DCC
R_{free} test set	48618 reflections (5.37%)	wwPDB-VP
Wilson B-factor (Å ²)	56.2	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 62.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7204	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/1577	0.41	0/2134
2	B	0.24	0/1558	0.40	0/2109
2	C	0.24	0/1554	0.41	0/2103
2	D	0.24	0/1568	0.40	0/2121
3	E	0.72	1/233 (0.4%)	0.74	0/360
3	H	0.71	1/233 (0.4%)	0.73	0/360
4	F	0.71	1/229 (0.4%)	0.72	0/353
4	G	0.71	1/229 (0.4%)	0.72	0/353
All	All	0.34	4/7181 (0.1%)	0.47	0/9893

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	1	A	OP3-P	-10.57	1.48	1.61
4	G	1	A	OP3-P	-10.56	1.48	1.61
3	E	1	A	OP3-P	-10.55	1.48	1.61
3	H	1	A	OP3-P	-10.56	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1537	0	1522	11	0
2	B	1525	0	1502	15	0
2	C	1522	0	1510	18	0
2	D	1536	0	1511	17	0
3	E	209	0	106	1	0
3	H	209	0	106	4	0
4	F	206	0	106	3	0
4	G	206	0	106	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	66	0	0	0	0
6	B	30	0	0	3	0
6	C	33	0	0	3	0
6	D	76	0	0	5	0
6	E	14	0	0	0	0
6	F	9	0	0	2	0
6	G	12	0	0	0	0
6	H	10	0	0	1	0
All	All	7204	0	6469	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:7:G:N7	6:H:101:HOH:O	2.24	0.70
2:C:4[B]:LEU:HD11	2:C:149:VAL:HG22	1.74	0.69
2:D:171[A]:GLN:NE2	6:D:402:HOH:O	2.26	0.68
1:A:32:TYR:OH	1:A:62:GLU:OE2	2.11	0.64
2:C:18:ARG:HH22	3:H:1:A:H2	1.46	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/215 (86%)	182 (98%)	4 (2%)	0	100	100
2	B	182/215 (85%)	178 (98%)	4 (2%)	0	100	100
2	C	182/215 (85%)	179 (98%)	3 (2%)	0	100	100
2	D	184/215 (86%)	180 (98%)	4 (2%)	0	100	100
All	All	734/860 (85%)	719 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/195 (88%)	170 (99%)	2 (1%)	67	75
2	B	169/194 (87%)	167 (99%)	2 (1%)	67	75
2	C	169/194 (87%)	169 (100%)	0	100	100
2	D	171/194 (88%)	169 (99%)	2 (1%)	67	75
All	All	681/777 (88%)	675 (99%)	6 (1%)	75	82

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	53	ASP
2	D	17	ARG
2	D	130	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	75	GLN
1	A	47	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	E	9/10 (90%)	1 (11%)	0
3	H	9/10 (90%)	1 (11%)	0
4	F	9/10 (90%)	1 (11%)	0
4	G	9/10 (90%)	1 (11%)	0
All	All	36/40 (90%)	4 (11%)	0

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	E	10	A
4	F	10	C
4	G	10	C
3	H	10	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSD	B	73	2	4,7,8	4.18	1 (25%)	1,8,10	3.15	1 (100%)
2	CSD	C	73	2	4,7,8	4.20	1 (25%)	1,8,10	2.46	1 (100%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSD	D	73	2	4,7,8	4.13	1 (25%)	1,8,10	3.21	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	B	73	2	-	1/2/6/8	-
2	CSD	C	73	2	-	2/2/6/8	-
2	CSD	D	73	2	-	1/2/6/8	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	73	CSD	OD1-SG	-8.15	1.40	1.47
2	B	73	CSD	OD1-SG	-8.09	1.40	1.47
2	D	73	CSD	OD1-SG	-8.01	1.40	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	73	CSD	OD1-SG-CB	3.21	111.52	105.60
2	B	73	CSD	OD1-SG-CB	3.15	111.40	105.60
2	C	73	CSD	OD1-SG-CB	2.46	110.14	105.60

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	73	CSD	CA-CB-SG-OD1
2	C	73	CSD	CA-CB-SG-OD1
2	D	73	CSD	CA-CB-SG-OD1
2	C	73	CSD	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	186/215 (86%)	-0.14	3 (1%) 70 71	31, 69, 119, 147	2 (1%)
2	B	181/215 (84%)	0.04	0 100 100	33, 91, 136, 171	3 (1%)
2	C	183/215 (85%)	0.12	4 (2%) 62 62	46, 92, 141, 175	1 (0%)
2	D	182/215 (84%)	-0.16	1 (0%) 87 89	27, 68, 108, 157	4 (2%)
3	E	10/10 (100%)	-0.33	0 100 100	56, 76, 88, 166	0
3	H	10/10 (100%)	-0.26	0 100 100	71, 79, 97, 132	0
4	F	10/10 (100%)	-0.22	0 100 100	70, 76, 93, 128	0
4	G	10/10 (100%)	-0.24	0 100 100	60, 76, 90, 125	0
All	All	772/900 (85%)	-0.05	8 (1%) 79 80	27, 80, 135, 175	10 (1%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	2	ALA	2.8
2	C	24	TYR	2.8
2	D	2	ALA	2.5
2	C	29	LEU	2.5
1	A	38	ASN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSD	B	73	8/9	0.77	0.13	102,120,152,155	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CSD	C	73	8/9	0.82	0.11	101,127,169,175	0
2	CSD	D	73	8/9	0.91	0.14	80,91,132,142	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ZN	C	301	1/1	0.98	0.09	91,91,91,91	0
5	ZN	A	300	1/1	0.99	0.13	71,71,71,71	0
5	ZN	D	301	1/1	0.99	0.13	68,68,68,68	0
5	ZN	B	301	1/1	1.00	0.06	81,81,81,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.