



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 16, 2024 – 06:13 AM EST

PDB ID : 5WGM  
Title : Crystal structure of Danio rerio histone deacetylase 6 catalytic domain 2 in complex with ACY-1083  
Authors : Porter, N.J.; Christianson, D.W.  
Deposited on : 2017-07-14  
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	 2% 95% 5%
1	B	364	 % 93% 7%
1	C	364	 2% 91% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	811	-	-	X	-
5	EDO	B	807	-	-	X	-
6	PGE	A	816	-	-	X	-
6	PGE	C	816	-	-	X	-

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9882 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hdac6 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	364	Total	C	N	O	S	0	12	0
			2904	1816	522	547	19			
1	B	363	Total	C	N	O	S	0	8	0
			2859	1793	510	537	19			
1	C	361	Total	C	N	O	S	0	11	0
			2867	1796	515	537	19			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	435	SER	-	expression tag	UNP A7YT55
A	436	ASN	-	expression tag	UNP A7YT55
A	437	ALA	-	expression tag	UNP A7YT55
A	438	GLY	-	expression tag	UNP A7YT55
A	439	GLY	-	expression tag	UNP A7YT55
B	435	SER	-	expression tag	UNP A7YT55
B	436	ASN	-	expression tag	UNP A7YT55
B	437	ALA	-	expression tag	UNP A7YT55
B	438	GLY	-	expression tag	UNP A7YT55
B	439	GLY	-	expression tag	UNP A7YT55
C	435	SER	-	expression tag	UNP A7YT55
C	436	ASN	-	expression tag	UNP A7YT55
C	437	ALA	-	expression tag	UNP A7YT55
C	438	GLY	-	expression tag	UNP A7YT55
C	439	GLY	-	expression tag	UNP A7YT55

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	B	2	Total	K	0	0
			2	2		
3	C	2	Total	K	0	0
			2	2		

- Molecule 4 is 2-[(4,4-difluoro-1-phenylcyclohexyl)amino]-N-hydroxypyrimidine-5-carboxamide (three-letter code: AH7) (formula: C<sub>17</sub>H<sub>18</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			25	17	2	4	2		
4	B	1	Total	C	F	N	O	0	0
			25	17	2	4	2		
4	C	1	Total	C	F	N	O	0	0
			25	17	2	4	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 8	C 4	O 4	0	1
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0
5	C	1	Total 8	C 4	O 4	0	1
5	C	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	4	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		
6	C	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	340	Total	O	0	5
			345	345		
8	B	311	Total	O	0	6
			317	317		
8	C	311	Total	O	0	5
			316	316		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hdac6 protein



- Molecule 1: Hdac6 protein



- Molecule 1: Hdac6 protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.81Å 174.30Å 149.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.95 – 1.75 49.95 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.95-1.75) 99.9 (49.95-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.26 (at 1.75Å)	Xtriage
Refinement program	PHENIX (dev_2776: ???)	Depositor
R, $R_{free}$	0.159 , 0.185 0.162 , 0.187	Depositor DCC
$R_{free}$ test set	6484 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.8	Xtriage
Anisotropy	0.396	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.028 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9882	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AH7, PEG, ZN, K, PGE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/2981	0.65	0/4042
1	B	0.56	0/2930	0.68	1/3976 (0.0%)
1	C	0.53	0/2939	0.66	0/3986
All	All	0.55	0/8850	0.66	1/12004 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	562	ARG	NE-CZ-NH2	5.25	122.93	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2904	0	2790	18	0
1	B	2859	0	2731	25	0
1	C	2867	0	2747	30	3
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	25	0	0	0	0
4	B	25	0	0	0	0
4	C	25	0	0	0	0
5	A	44	0	66	13	3
5	B	64	0	96	17	0
5	C	48	0	72	9	0
6	A	10	0	14	6	0
6	C	10	0	14	7	0
7	A	7	0	9	0	0
7	C	7	0	10	3	0
8	A	345	0	0	5	1
8	B	317	0	0	4	1
8	C	316	0	0	4	1
All	All	9882	0	8549	78	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:485:ARG:HH21	6:C:816:PGE:H42	1.31	0.95
1:C:762:LEU:HD23	6:C:816:PGE:H1	1.53	0.90
1:A:488:ARG:HH12	5:A:805:EDO:H21	1.40	0.85
1:C:485:ARG:HE	6:C:816:PGE:H5	1.43	0.84
6:A:816:PGE:H52	1:B:746:ASN:ND2	1.99	0.77

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:497:GLU:CD	5:A:811:EDO:O1[5_545]	1.38	0.82
1:C:497:GLU:OE2	5:A:811:EDO:O1[5_545]	1.73	0.47
1:C:497:GLU:OE1	5:A:811:EDO:O1[5_545]	1.80	0.40
8:A:1208:HOH:O	8:B:1178:HOH:O[3_554]	2.14	0.06
8:C:997:HOH:O	8:C:997:HOH:O[4_555]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/364 (103%)	366 (98%)	8 (2%)	0	100	100
1	B	369/364 (101%)	362 (98%)	7 (2%)	0	100	100
1	C	368/364 (101%)	361 (98%)	7 (2%)	0	100	100
All	All	1111/1092 (102%)	1089 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/305 (103%)	309 (99%)	4 (1%)	65	51
1	B	304/305 (100%)	297 (98%)	7 (2%)	45	25
1	C	306/305 (100%)	301 (98%)	5 (2%)	58	42
All	All	923/915 (101%)	907 (98%)	16 (2%)	54	39

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	573	HIS
1	C	530	ASN
1	B	611	TRP
1	C	465	GLU
1	B	573	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 55 ligands modelled in this entry, 9 are monoatomic - leaving 46 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	EDO	A	812	-	3,3,3	0.35	0	2,2,2	0.51	0
6	PGE	A	816	-	9,9,9	0.36	0	8,8,8	0.61	0
5	EDO	C	814	-	3,3,3	0.46	0	2,2,2	0.42	0
5	EDO	A	805	-	3,3,3	0.42	0	2,2,2	0.38	0
5	EDO	C	810	-	3,3,3	0.70	0	2,2,2	0.54	0
5	EDO	C	805	-	3,3,3	0.42	0	2,2,2	0.48	0
5	EDO	C	812	-	3,3,3	0.37	0	2,2,2	0.28	0
5	EDO	A	810	-	3,3,3	0.21	0	2,2,2	1.41	0
5	EDO	C	808	-	3,3,3	0.44	0	2,2,2	0.25	0
4	AH7	A	804	2	25,27,27	1.60	5 (20%)	31,39,39	1.73	8 (25%)
5	EDO	A	809	-	3,3,3	0.16	0	2,2,2	0.95	0
5	EDO	A	807	-	3,3,3	0.55	0	2,2,2	0.28	0
5	EDO	B	812	-	3,3,3	0.53	0	2,2,2	0.10	0
5	EDO	C	807	-	3,3,3	0.39	0	2,2,2	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	A	815	-	3,3,3	0.67	0	2,2,2	0.12	0
5	EDO	B	810	-	3,3,3	0.37	0	2,2,2	0.52	0
5	EDO	B	805	-	3,3,3	0.57	0	2,2,2	0.36	0
4	AH7	B	804	2	25,27,27	2.21	9 (36%)	31,39,39	1.72	5 (16%)
4	AH7	C	804	2	25,27,27	1.74	5 (20%)	31,39,39	1.46	4 (12%)
5	EDO	B	814	-	3,3,3	0.45	0	2,2,2	0.01	0
7	PEG	A	817	-	6,6,6	0.97	0	5,5,5	0.86	0
5	EDO	C	815	-	3,3,3	0.45	0	2,2,2	0.37	0
5	EDO	B	811	-	3,3,3	0.52	0	2,2,2	0.34	0
5	EDO	B	807	-	3,3,3	0.42	0	2,2,2	0.24	0
5	EDO	A	813	-	3,3,3	0.56	0	2,2,2	0.69	0
5	EDO	B	813	-	3,3,3	0.35	0	2,2,2	0.48	0
5	EDO	C	809[B]	-	3,3,3	0.55	0	2,2,2	0.31	0
5	EDO	B	818	-	3,3,3	0.42	0	2,2,2	0.53	0
5	EDO	B	806	-	3,3,3	0.24	0	2,2,2	0.80	0
5	EDO	C	806	-	3,3,3	0.37	0	2,2,2	0.73	0
5	EDO	B	808	-	3,3,3	0.43	0	2,2,2	0.32	0
5	EDO	C	809[A]	-	3,3,3	0.48	0	2,2,2	0.20	0
5	EDO	A	814	-	3,3,3	0.39	0	2,2,2	0.41	0
5	EDO	B	816[B]	-	3,3,3	0.55	0	2,2,2	0.46	0
5	EDO	B	819	-	3,3,3	0.53	0	2,2,2	0.29	0
5	EDO	B	809	-	3,3,3	0.41	0	2,2,2	0.41	0
6	PGE	C	816	-	9,9,9	0.38	0	8,8,8	0.55	0
5	EDO	B	816[A]	-	3,3,3	0.44	0	2,2,2	0.43	0
5	EDO	B	817	-	3,3,3	0.45	0	2,2,2	0.31	0
5	EDO	B	815	-	3,3,3	0.40	0	2,2,2	0.53	0
5	EDO	A	811	-	3,3,3	0.58	0	2,2,2	0.45	0
5	EDO	C	811	-	3,3,3	0.51	0	2,2,2	0.33	0
5	EDO	C	813	-	3,3,3	0.43	0	2,2,2	0.22	0
7	PEG	C	817	-	6,6,6	0.75	0	5,5,5	0.58	0
5	EDO	A	806	-	3,3,3	0.53	0	2,2,2	0.47	0
5	EDO	A	808	-	3,3,3	0.40	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	812	-	-	0/1/1/1	-
6	PGE	A	816	-	-	6/7/7/7	-
5	EDO	C	814	-	-	1/1/1/1	-

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	805	-	-	1/1/1/1	-
5	EDO	C	810	-	-	1/1/1/1	-
5	EDO	C	805	-	-	0/1/1/1	-
5	EDO	C	812	-	-	1/1/1/1	-
5	EDO	A	810	-	-	0/1/1/1	-
5	EDO	C	808	-	-	1/1/1/1	-
4	AH7	A	804	2	-	3/17/31/31	0/3/3/3
5	EDO	A	809	-	-	1/1/1/1	-
5	EDO	A	807	-	-	1/1/1/1	-
5	EDO	B	812	-	-	1/1/1/1	-
5	EDO	C	807	-	-	0/1/1/1	-
5	EDO	A	815	-	-	1/1/1/1	-
5	EDO	B	810	-	-	0/1/1/1	-
5	EDO	B	805	-	-	0/1/1/1	-
4	AH7	B	804	2	-	3/17/31/31	0/3/3/3
4	AH7	C	804	2	-	3/17/31/31	0/3/3/3
5	EDO	B	814	-	-	0/1/1/1	-
7	PEG	A	817	-	-	4/4/4/4	-
5	EDO	C	815	-	-	0/1/1/1	-
5	EDO	B	811	-	-	1/1/1/1	-
5	EDO	B	807	-	-	1/1/1/1	-
5	EDO	A	813	-	-	1/1/1/1	-
5	EDO	B	813	-	-	0/1/1/1	-
5	EDO	C	809[B]	-	-	0/1/1/1	-
5	EDO	B	818	-	-	1/1/1/1	-
5	EDO	B	806	-	-	1/1/1/1	-
5	EDO	C	806	-	-	0/1/1/1	-
5	EDO	B	808	-	-	1/1/1/1	-
5	EDO	C	809[A]	-	-	1/1/1/1	-
5	EDO	A	814	-	-	0/1/1/1	-
5	EDO	B	816[B]	-	-	1/1/1/1	-
5	EDO	B	819	-	-	0/1/1/1	-
5	EDO	B	809	-	-	0/1/1/1	-
6	PGE	C	816	-	-	4/7/7/7	-
5	EDO	B	816[A]	-	-	0/1/1/1	-
5	EDO	B	817	-	-	1/1/1/1	-
5	EDO	B	815	-	-	1/1/1/1	-
5	EDO	A	811	-	-	0/1/1/1	-
5	EDO	C	811	-	-	0/1/1/1	-
5	EDO	C	813	-	-	1/1/1/1	-
7	PEG	C	817	-	-	2/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	806	-	-	1/1/1/1	-
5	EDO	A	808	-	-	1/1/1/1	-

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	804	AH7	C10-C11	-5.40	1.46	1.53
4	B	804	AH7	C08-N09	4.56	1.41	1.34
4	C	804	AH7	C08-N09	3.77	1.40	1.34
4	A	804	AH7	C10-C11	-3.42	1.49	1.53
4	B	804	AH7	C08-N07	3.23	1.39	1.34

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	804	AH7	C22-C19-C18	5.06	119.36	114.33
4	A	804	AH7	N24-C08-N07	-4.88	121.49	126.43
4	B	804	AH7	N24-C08-N07	-4.74	121.63	126.43
4	C	804	AH7	N24-C08-N07	-4.60	121.77	126.43
4	A	804	AH7	F20-C19-C22	3.30	114.60	108.92

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	804	AH7	C23-C10-C11-C16
4	B	804	AH7	C23-C10-C11-C16
4	C	804	AH7	C23-C10-C11-C16
6	C	816	PGE	O3-C5-C6-O4
6	A	816	PGE	O3-C5-C6-O4

There are no ring outliers.

25 monomers are involved in 58 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	816	PGE	6	0
5	C	814	EDO	1	0
5	A	805	EDO	3	0
5	C	810	EDO	1	0
5	A	810	EDO	1	0
5	A	809	EDO	2	0


*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	812	EDO	1	0
5	C	807	EDO	2	0
5	A	815	EDO	1	0
5	B	810	EDO	3	0
5	B	807	EDO	4	0
5	B	813	EDO	1	0
5	C	809[B]	EDO	1	0
5	B	818	EDO	2	0
5	C	806	EDO	3	0
5	A	814	EDO	1	0
5	B	816[B]	EDO	3	0
5	B	809	EDO	1	0
6	C	816	PGE	7	0
5	B	816[A]	EDO	2	0
5	A	811	EDO	1	3
5	C	813	EDO	1	0
7	C	817	PEG	3	0
5	A	806	EDO	3	0
5	A	808	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

## Ligand AH7 A 804



## Ligand AH7 B 804





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	364/364 (100%)	-0.47	7 (1%) 66 72	4, 14, 27, 51	13 (3%)
1	B	363/364 (99%)	-0.44	5 (1%) 73 79	4, 14, 28, 58	10 (2%)
1	C	361/364 (99%)	-0.42	8 (2%) 62 68	4, 14, 28, 52	12 (3%)
All	All	1088/1092 (99%)	-0.44	20 (1%) 67 73	4, 14, 28, 58	35 (3%)

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	772	LEU	5.6
1	C	773	THR	4.8
1	A	435	SER	4.1
1	B	436	ASN	4.0
1	C	436	ASN	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

## 6.5 Other polymers [i](#)

There are no such residues in this entry.