



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 09:29 PM EDT

PDB ID : 3WPG  
Title : Crystal structure of mouse TLR9 in complex with inhibitory DNA4084 (form 1)  
Authors : Ohto, U.; Shimizu, T.  
Deposited on : 2014-01-11  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3139 (2.26-2.22)
Clashscore	180529	3381 (2.26-2.22)
Ramachandran outliers	177936	3334 (2.26-2.22)
Sidechain outliers	177891	3335 (2.26-2.22)
RSRZ outliers	164620	3138 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	803	
2	B	12	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	S	11	0	0
			5757	3676	1015	1037	29			

There are 17 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	22	ARG	-	expression tag	UNP Q9EQU3
A	23	SER	-	expression tag	UNP Q9EQU3
A	24	PRO	-	expression tag	UNP Q9EQU3
A	25	TRP	-	expression tag	UNP Q9EQU3
A	200	GLN	ASN	engineered mutation	UNP Q9EQU3
A	242	GLN	ASN	engineered mutation	UNP Q9EQU3
A	309	GLN	ASN	engineered mutation	UNP Q9EQU3
A	495	GLN	ASN	engineered mutation	UNP Q9EQU3
A	568	GLN	ASN	engineered mutation	UNP Q9EQU3
A	695	GLN	ASN	engineered mutation	UNP Q9EQU3
A	752	GLN	ASN	engineered mutation	UNP Q9EQU3
A	819	GLU	-	expression tag	UNP Q9EQU3
A	820	PHE	-	expression tag	UNP Q9EQU3
A	821	LEU	-	expression tag	UNP Q9EQU3
A	822	VAL	-	expression tag	UNP Q9EQU3
A	823	PRO	-	expression tag	UNP Q9EQU3
A	824	ARG	-	expression tag	UNP Q9EQU3

- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*CP\*TP\*GP\*GP\*AP\*TP\*GP\*GP\*GP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	P	0	0	0
			227	108	45	64	10			


- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:

C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	68	Total	O	0	0
			68	68		
5	B	9	Total	O	0	0
			9	9		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Toll-like receptor 9



#### • Molecule 2: DNA (5'-D(\*CP\*CP\*TP\*GP\*GP\*AP\*TP\*GP\*GP\*GP\*AP\*A)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.23Å 123.86Å 130.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.62 – 2.25 35.62 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.62-2.25) 99.9 (35.62-2.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.8.1 _1168	Depositor
R, $R_{free}$	0.236 , 0.280 0.240 , 0.282	Depositor DCC
$R_{free}$ test set	2825 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtriage
Anisotropy	0.877	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 55.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6100	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.26	0/5878	0.46	0/7972
2	B	0.46	0/255	1.10	1/393 (0.3%)
All	All	0.27	0/6133	0.51	1/8365 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	DG	O4'-C1'-N9	6.67	112.67	108.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5757	0	5811	70	0
2	B	227	0	125	1	0
3	A	14	0	13	0	0
4	A	20	0	0	1	0
4	B	5	0	0	0	0
5	A	68	0	0	0	1
5	B	9	0	0	0	0
All	All	6100	0	5949	71	1



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:ASN:HD21	1:A:397:HIS:HB3	1.35	0.91
1:A:749:ILE:HG23	1:A:752:GLN:HE21	1.56	0.70
1:A:214:VAL:O	1:A:216:ARG:NH1	2.26	0.69
1:A:426:ARG:CZ	1:A:484:ASN:HD21	2.10	0.65
1:A:300:ASN:OD1	1:A:302:SER:OG	2.14	0.64
1:A:372:ASN:HD21	1:A:397:HIS:CB	2.09	0.64
1:A:117:ARG:HA	1:A:120:LEU:HD23	1.82	0.62
1:A:51:LYS:N	4:A:904:SO4:O4	2.28	0.61
1:A:737:ILE:HA	1:A:762:PRO:HD2	1.83	0.61
1:A:215:PRO:HB2	1:A:218:LEU:HD21	1.84	0.59
1:A:558:GLN:H	1:A:587:ASP:HB2	1.67	0.59
1:A:372:ASN:ND2	1:A:397:HIS:HB3	2.12	0.59
1:A:522:THR:OG1	1:A:523:ASN:OD1	2.20	0.58
1:A:564:GLY:O	1:A:591:ARG:NH1	2.37	0.58
1:A:325:THR:OG1	1:A:355:HIS:O	2.16	0.57
1:A:493:PHE:HA	1:A:496:LEU:HD12	1.88	0.56
1:A:129:ASN:OD1	1:A:131:SER:OG	2.16	0.55
1:A:706:LYS:HG2	1:A:730:GLU:HB2	1.89	0.54
1:A:568:GLN:HB2	1:A:592:VAL:HA	1.90	0.54
1:A:740:THR:OG1	1:A:741:VAL:N	2.40	0.53
1:A:554:SER:OG	1:A:582:SER:OG	2.23	0.53
1:A:370:ASN:HA	1:A:397:HIS:HB2	1.91	0.52
1:A:787:ASN:O	1:A:790:LYS:NZ	2.31	0.52
1:A:353:ARG:HD3	1:A:380:ASN:ND2	2.24	0.52
1:A:741:VAL:HG22	1:A:774:LEU:HD23	1.90	0.52
1:A:668:PHE:HB3	1:A:695:GLN:HE22	1.75	0.52
1:A:785:LEU:HD12	1:A:789:VAL:HG11	1.91	0.52
1:A:304:PHE:HB3	1:A:333:LEU:HD21	1.91	0.52
1:A:573:THR:HB	1:A:597:ASN:HB2	1.92	0.51
1:A:646:PRO:HB3	1:A:676:PHE:HE2	1.75	0.50
1:A:50:LEU:HD13	1:A:54:PRO:HG3	1.93	0.49
1:A:741:VAL:HG13	1:A:771:PHE:HB2	1.94	0.49
1:A:757:ASP:OD1	1:A:759:ARG:HD3	2.13	0.49
1:A:336:LEU:HD21	1:A:339:LEU:HD13	1.96	0.48
1:A:370:ASN:N	1:A:370:ASN:OD1	2.48	0.47
1:A:746:PHE:CB	1:A:750:VAL:HB	2.45	0.47
1:A:575:LEU:HD12	1:A:578:LEU:HD22	1.97	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:LEU:HG	1:A:387:LEU:HD12	1.97	0.47
1:A:668:PHE:HB3	1:A:695:GLN:NE2	2.30	0.46
1:A:339:LEU:HG	1:A:341:LEU:HD13	1.96	0.46
1:A:250:ASP:HA	1:A:290:VAL:HG13	1.95	0.46
1:A:751:MET:HE3	1:A:777:GLU:CD	2.36	0.46
1:A:247:ARG:NH1	1:A:287:GLU:OE2	2.48	0.45
1:A:163:LEU:HB3	1:A:166:LEU:HD12	1.99	0.45
1:A:220:PRO:HB3	1:A:242:GLN:HG3	1.99	0.45
1:A:283:LEU:HD23	1:A:286:LEU:HD13	1.99	0.45
1:A:42:LEU:HA	1:A:67:ARG:HB2	1.98	0.45
1:A:759:ARG:HD2	1:A:783:PRO:O	2.16	0.45
1:A:693:LEU:HB2	1:A:717:VAL:HB	1.99	0.44
1:A:207:LYS:HD2	1:A:228:SER:OG	2.17	0.44
1:A:646:PRO:HB3	1:A:676:PHE:CE2	2.52	0.44
2:B:4:DG:N3	2:B:4:DG:H5"	2.33	0.44
1:A:86:LEU:HB3	1:A:89:LEU:HG	1.99	0.44
1:A:698:LEU:O	1:A:724:LEU:HB2	2.17	0.44
1:A:353:ARG:HD3	1:A:380:ASN:HD21	1.83	0.43
1:A:101:THR:HA	1:A:102:GLY:HA2	1.59	0.43
1:A:227:VAL:O	1:A:230:ASN:ND2	2.49	0.43
1:A:695:GLN:N	1:A:696:GLY:HA3	2.33	0.43
1:A:380:ASN:ND2	1:A:383:THR:OG1	2.52	0.43
1:A:337:ARG:NH1	1:A:367:GLN:HE21	2.16	0.43
1:A:693:LEU:O	1:A:718:VAL:HG12	2.17	0.43
1:A:749:ILE:O	1:A:752:GLN:HG2	2.19	0.42
1:A:680:LEU:HD21	1:A:683:LEU:HB2	2.01	0.42
1:A:672:THR:HG23	1:A:699:PRO:HB3	2.02	0.42
1:A:83:PHE:HD1	1:A:86:LEU:HD12	1.84	0.42
1:A:261:ALA:HA	1:A:262:PRO:HD3	1.84	0.41
1:A:337:ARG:NH1	1:A:365:SER:O	2.52	0.41
1:A:166:LEU:O	1:A:198:LEU:HD12	2.20	0.41
1:A:201:LEU:HD12	1:A:201:LEU:HA	1.94	0.41
1:A:228:SER:HB3	1:A:250:ASP:OD1	2.21	0.41
1:A:741:VAL:HA	1:A:745:TRP:CH2	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1024:HOH:O	5:A:1033:HOH:O[2_555]	2.12	0.08

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	714/803 (89%)	661 (93%)	49 (7%)	4 (1%)	22	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	476	PHE
1	A	690	LEU
1	A	383	THR
1	A	769	ALA

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	662/728 (91%)	615 (93%)	47 (7%)	12	9

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	THR
1	A	43	VAL
1	A	62	CYS
1	A	65	ILE
1	A	76	HIS
1	A	81	SER
1	A	96	TRP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	138	VAL
1	A	156	LEU
1	A	169	LEU
1	A	184	CYS
1	A	198	LEU
1	A	200	GLN
1	A	248	VAL
1	A	272	SER
1	A	273	LEU
1	A	290	VAL
1	A	302	SER
1	A	309	GLN
1	A	313	LEU
1	A	325	THR
1	A	335	ARG
1	A	354	LEU
1	A	370	ASN
1	A	372	ASN
1	A	385	ARG
1	A	396	LEU
1	A	418	ARG
1	A	420	VAL
1	A	565	ILE
1	A	573	THR
1	A	576	SER
1	A	579	GLN
1	A	588	ILE
1	A	629	SER
1	A	667	SER
1	A	672	THR
1	A	693	LEU
1	A	695	GLN
1	A	715	VAL
1	A	717	VAL
1	A	726	VAL
1	A	731	VAL
1	A	741	VAL
1	A	753	LEU
1	A	755	VAL
1	A	776	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	372	ASN
1	A	484	ASN
1	A	752	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	901	1	14,14,15	0.46	0	17,19,21	0.81	1 (5%)
4	SO4	A	904	-	4,4,4	0.24	0	6,6,6	0.12	0
4	SO4	A	903	-	4,4,4	0.23	0	6,6,6	0.07	0
4	SO4	B	101	-	4,4,4	0.23	0	6,6,6	0.08	0
4	SO4	A	902	-	4,4,4	0.22	0	6,6,6	0.06	0
4	SO4	A	905	-	4,4,4	0.23	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '–' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	901	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	NAG	C1-O5-C5	2.46	115.48	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	904	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	728/803 (90%)	0.73	68 (9%) 16 15	15, 55, 90, 110	2 (0%)
2	B	11/12 (91%)	-0.03	0 100 100	32, 43, 51, 80	0
All	All	739/815 (90%)	0.72	68 (9%) 16 16	15, 54, 90, 110	2 (0%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281	HIS	5.9
1	A	102	GLY	5.1
1	A	352	ALA	5.0
1	A	695	GLN	4.9
1	A	573	THR	4.7
1	A	588	ILE	4.6
1	A	109	SER	4.5
1	A	474	PHE	4.4
1	A	60	ALA	4.3
1	A	767	CYS	3.8
1	A	101	THR	3.5
1	A	28	THR	3.4
1	A	47	TRP	3.3
1	A	346	ARG	3.2
1	A	470	ARG	3.2
1	A	400	MET	3.2
1	A	308	VAL	3.1
1	A	558	GLN	3.1
1	A	589	HIS	3.0
1	A	697	THR	3.0
1	A	266	ILE	3.0
1	A	377	ARG	3.0
1	A	196	LEU	2.9
1	A	305	GLN	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	537	TYR	2.8
1	A	513	VAL	2.8
1	A	347	LYS	2.8
1	A	259	ASP	2.8
1	A	332	ASN	2.7
1	A	468	MET	2.7
1	A	62	CYS	2.7
1	A	280	PHE	2.7
1	A	41	GLY	2.6
1	A	748	PRO	2.6
1	A	276	HIS	2.5
1	A	432	THR	2.5
1	A	90	ARG	2.5
1	A	306	GLY	2.5
1	A	564	GLY	2.5
1	A	772	VAL	2.5
1	A	694	THR	2.5
1	A	63	SER	2.5
1	A	376	PHE	2.5
1	A	324	ILE	2.4
1	A	277	PRO	2.4
1	A	224	TYR	2.4
1	A	124	THR	2.4
1	A	565	ILE	2.4
1	A	405	GLN	2.4
1	A	613	ARG	2.3
1	A	696	GLY	2.3
1	A	776	LEU	2.3
1	A	375	PHE	2.3
1	A	262	PRO	2.2
1	A	587	ASP	2.2
1	A	809	LEU	2.2
1	A	431	SER	2.2
1	A	469	ASP	2.2
1	A	96	TRP	2.2
1	A	608	GLY	2.1
1	A	701	GLY	2.1
1	A	720	ALA	2.1
1	A	554	SER	2.1
1	A	257	ARG	2.1
1	A	37	LEU	2.1
1	A	242	GLN	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	325	THR	2.0
1	A	381	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	A	902	5/5	0.60	0.17	138,139,140,142	0
4	SO4	A	903	5/5	0.74	0.22	119,119,119,121	0
4	SO4	A	905	5/5	0.77	0.09	102,103,106,106	0
4	SO4	B	101	5/5	0.81	0.23	103,107,107,110	0
3	NAG	A	901	14/15	0.84	0.11	45,56,63,65	0
4	SO4	A	904	5/5	0.85	0.13	110,111,113,114	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.