



wwPDB NMR Structure Validation Summary Report ⓘ

Dec 25, 2024 – 06:59 PM EST

PDB ID : 7X7S
BMRB ID : 18133
Title : Solution structure of human adenylate kinase 1 (hAK1)
Authors : Zhang, H.
Deposited on : 2022-03-10

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment is 85%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	199	

2 Ensemble composition and analysis

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:199 (194)	0.71	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 8, 11, 12, 13, 15, 16, 17, 19
2	2, 5, 6, 7, 10, 18, 20
3	9, 14

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3078 atoms, of which 1561 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Adenylate kinase isoenzyme 1.

Mol	Chain	Residues	Atoms						Trace
1	A	194	Total	C	H	N	O	S	0
			3078	951	1561	264	295	7	

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P00568
A	2	PRO	-	expression tag	UNP P00568
A	3	LEU	-	expression tag	UNP P00568
A	4	GLY	-	expression tag	UNP P00568
A	5	SER	-	expression tag	UNP P00568

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Adenylate kinase isoenzyme 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 300 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA2alpha	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	2300
Number of shifts mapped to atoms	2300
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	85%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1517	1561	1558	136±9
All	All	30340	31220	31160	2716

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 44.

5 of 562 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:47:LEU:HD21	1:A:84:ALA:HB2	0.89	1.45	2	3
1:A:74:LEU:H	1:A:74:LEU:HD12	0.86	1.27	19	5
1:A:131:THR:OG1	1:A:151:ILE:HG21	0.83	1.73	3	19
1:A:74:LEU:HD12	1:A:74:LEU:N	0.82	1.88	5	17
1:A:198:LEU:HD13	1:A:199:LYS:N	0.81	1.90	7	17

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/199 (96%)	168±2 (87±1%)	21±2 (11±1%)	4±1 (2±1%)	8	50
All	All	3840/3980 (96%)	3350 (87%)	413 (11%)	77 (2%)	8	50

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	101	PRO	20
1	A	127	PRO	20
1	A	198	LEU	8
1	A	143	ARG	7
1	A	144	VAL	6

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/170 (98%)	123±3 (74±2%)	44±3 (26±2%)	2	21
All	All	3340/3400 (98%)	2462 (74%)	878 (26%)	2	21

5 of 91 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	12	LYS	20
1	A	15	ILE	20
1	A	17	PHE	20
1	A	18	VAL	20
1	A	37	TYR	20

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 85% for the well-defined parts and 85% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: `cs_list`

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2300
Number of shifts mapped to atoms	2300
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	16

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	188	-1.58 ± 0.12	Should be checked
$^{13}\text{C}_\beta$	163	-0.99 ± 0.10	Should be checked
$^{13}\text{C}'$	189	-1.48 ± 0.15	Should be applied
^{15}N	186	-0.65 ± 0.17	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 85%, i.e. 2258 atoms were assigned a chemical shift out of a possible 2664. 0 out of 35 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	958/977 (98%)	395/401 (99%)	377/388 (97%)	186/188 (99%)
Sidechain	1254/1558 (80%)	868/1007 (86%)	386/485 (80%)	0/66 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	46/129 (36%)	46/61 (75%)	0/64 (0%)	0/4 (0%)
Overall	2258/2664 (85%)	1309/1469 (89%)	763/937 (81%)	186/258 (72%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	13	THR	HG1	5.33	0.08 – 2.19	19.9
1	A	44	THR	HG1	4.76	0.08 – 2.19	17.2
1	A	140	THR	HG1	4.76	0.08 – 2.19	17.2
1	A	28	THR	HG1	4.36	0.08 – 2.19	15.3
1	A	157	THR	HG1	4.34	0.08 – 2.19	15.2
1	A	162	THR	HG1	4.04	0.08 – 2.19	13.8
1	A	32	LYS	HE3	4.62	1.92 – 3.89	8.7
1	A	49	ARG	HD3	1.11	1.81 – 4.39	-7.7
1	A	122	TYR	HE1	8.42	5.59 – 7.82	7.7
1	A	54	SER	HB2	2.13	2.61 – 5.13	-6.9
1	A	137	ARG	HH21	9.31	4.81 – 8.80	6.3
1	A	177	LYS	HE3	4.13	1.92 – 3.89	6.2
1	A	176	ARG	HH11	9.40	4.72 – 9.08	5.7
1	A	102	ARG	HD3	4.53	1.81 – 4.39	5.5
1	A	60	LYS	HE3	3.95	1.92 – 3.89	5.3
1	A	190	GLN	HB2	3.32	0.80 – 3.29	5.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	3005
Intra-residue ($ i-j =0$)	972
Sequential ($ i-j =1$)	778
Medium range ($ i-j >1$ and $ i-j <5$)	598
Long range ($ i-j \geq 5$)	657
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	350
Number of unmapped restraints	0
Number of restraints per residue	16.9
Number of long range restraints per residue ¹	3.3

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	119.5	0.2
0.2-0.5 (Medium)	122.2	0.5
>0.5 (Large)	210.4	5.21

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	50.1	6.62
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restraints type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	972	32.3	185	19.0	6.2	33	3.4	1.1
Backbone-Backbone	139	4.6	1	0.7	0.0	1	0.7	0.0
Backbone-Sidechain	614	20.4	147	23.9	4.9	23	3.7	0.8
Sidechain-Sidechain	219	7.3	37	16.9	1.2	9	4.1	0.3
Sequential ($i-j =1$)	778	25.9	233	29.9	7.8	35	4.5	1.2
Backbone-Backbone	270	9.0	20	7.4	0.7	7	2.6	0.2
Backbone-Sidechain	407	13.5	159	39.1	5.3	19	4.7	0.6
Sidechain-Sidechain	101	3.4	54	53.5	1.8	9	8.9	0.3
Medium range ($i-j >1$ & $i-j <5$)	598	19.9	215	36.0	7.2	49	8.2	1.6
Backbone-Backbone	172	5.7	24	14.0	0.8	3	1.7	0.1
Backbone-Sidechain	286	9.5	129	45.1	4.3	29	10.1	1.0
Sidechain-Sidechain	140	4.7	62	44.3	2.1	17	12.1	0.6
Long range ($i-j \geq 5$)	657	21.9	273	41.6	9.1	72	11.0	2.4
Backbone-Backbone	57	1.9	13	22.8	0.4	6	10.5	0.2
Backbone-Sidechain	317	10.5	137	43.2	4.6	35	11.0	1.2
Sidechain-Sidechain	283	9.4	123	43.5	4.1	31	11.0	1.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	3005	100.0	906	30.1	30.1	189	6.3	6.3
Backbone-Backbone	638	21.2	58	9.1	1.9	17	2.7	0.6
Backbone-Sidechain	1624	54.0	572	35.2	19.0	106	6.5	3.5
Sidechain-Sidechain	743	24.7	276	37.1	9.2	66	8.9	2.2

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	94	101	120	145	0	460	0.6	2.95	0.51	0.44
2	85	114	109	138	0	446	0.63	3.82	0.54	0.47
3	89	112	117	149	0	467	0.6	3.55	0.53	0.42
4	94	110	120	146	0	470	0.61	3.82	0.51	0.48
5	88	113	116	152	0	469	0.64	5.21	0.55	0.51
6	91	113	114	151	0	469	0.62	3.96	0.54	0.45
7	89	96	114	151	0	450	0.56	2.37	0.46	0.43
8	88	102	113	144	0	447	0.55	2.27	0.45	0.4
9	94	99	115	145	0	453	0.6	3.38	0.52	0.45
10	87	108	110	159	0	464	0.61	4.13	0.54	0.44

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	78	111	115	144	0	448	0.6	3.01	0.52	0.42
12	92	107	111	151	0	461	0.62	3.98	0.53	0.46
13	99	108	116	141	0	464	0.57	3.25	0.49	0.4
14	94	118	121	147	0	480	0.64	3.97	0.58	0.45
15	94	109	112	148	0	463	0.58	3.3	0.5	0.41
16	94	110	115	143	0	462	0.63	3.31	0.53	0.47
17	86	101	116	151	0	454	0.58	3.03	0.52	0.4
18	89	104	113	149	0	455	0.59	2.97	0.51	0.42
19	94	109	117	148	0	468	0.61	3.81	0.53	0.43
20	94	104	123	149	0	470	0.61	2.88	0.51	0.45

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2099(IR:787, SQ:545, MR:383, LR:384, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
31	39	37	42	0	149	1	5.0
19	22	21	20	0	82	2	10.0
9	12	10	11	0	42	3	15.0
6	10	4	11	0	31	4	20.0
10	14	10	13	0	47	5	25.0
6	11	7	7	0	31	6	30.0
6	8	5	7	0	26	7	35.0
8	7	4	9	0	28	8	40.0
2	9	6	5	0	22	9	45.0
6	6	2	6	0	20	10	50.0
6	12	11	8	0	37	11	55.0
6	4	4	5	0	19	12	60.0
5	2	2	5	0	14	13	65.0
7	9	8	8	0	32	14	70.0
6	9	8	4	0	27	15	75.0
3	3	4	6	0	16	16	80.0
3	3	2	5	0	13	17	85.0
6	7	13	8	0	34	18	90.0
7	11	8	21	0	47	19	95.0
33	35	49	72	0	189	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,573)	1:73:A:PRO:HD2	1:61:A:LYS:HE3	20	1.93	0.57	2.03
(1,864)	1:124:A:ASP:HB3	1:177:A:LYS:HD2	20	1.92	0.67	1.68
(1,1002)	1:155:A:LEU:HB3	1:154:A:ARG:HB2	20	1.86	0.29	1.9
(4,160)	1:97:A:ILE:HG21	1:110:A:PHE:HB2	20	1.83	0.1	1.84
(4,160)	1:97:A:ILE:HG21	1:77:A:VAL:HA	20	1.83	0.1	1.84
(1,599)	1:77:A:VAL:HG21	1:73:A:PRO:HD3	20	1.77	0.8	1.58
(4,219)	1:65:A:ILE:HA	1:71:A:LEU:HD11	20	1.74	0.94	1.9
(4,219)	1:65:A:ILE:HA	1:71:A:LEU:HD21	20	1.74	0.94	1.9
(4,219)	1:65:A:ILE:HA	1:72:A:VAL:HG22	20	1.74	0.94	1.9
(4,219)	1:65:A:ILE:HA	1:71:A:LEU:HD23	20	1.74	0.94	1.9
(4,219)	1:65:A:ILE:HA	1:72:A:VAL:HG21	20	1.74	0.94	1.9
(4,219)	1:65:A:ILE:HA	1:71:A:LEU:HD22	20	1.74	0.94	1.9
(1,781)	1:113:A:ARG:HD2	1:109:A:GLU:HB2	20	1.7	0.26	1.58
(4,147)	1:197:A:ALA:HB1	1:16:A:ILE:HD13	20	1.7	0.24	1.77
(4,147)	1:84:A:ALA:HB3	1:15:A:ILE:HD12	20	1.7	0.24	1.77
(4,147)	1:84:A:ALA:HB1	1:15:A:ILE:HD12	20	1.7	0.24	1.77

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(4,147)	1:197:A:ALA:HB1	1:121:A:LEU:HD21	20	1.7	0.24	1.77
(1,359)	1:28:A:THR:HA	1:31:A:GLU:HB3	20	1.67	0.04	1.67
(4,288)	1:93:A:LYS:H	1:16:A:ILE:HG13	20	1.61	0.07	1.6

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,599)	1:77:A:VAL:HG21	1:73:A:PRO:HD3	5	5.21
(1,53)	1:185:A:ASP:HB3	1:184:A:VAL:HG22	10	4.13

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,238)	1:9:A:LYS:HG2	1:10:A:LEU:HD11	12	3.98
(1,53)	1:185:A:ASP:HB3	1:184:A:VAL:HG22	14	3.97
(1,1228)	1:197:A:ALA:HA	1:198:A:LEU:HD11	6	3.96
(1,95)	1:184:A:VAL:HG11	1:25:A:GLY:HA2	14	3.86
(1,1350)	1:55:A:GLY:H	1:62:A:LEU:HD11	4	3.82
(1,251)	1:10:A:LEU:HD21	1:115:A:GLY:HA2	2	3.82
(1,1228)	1:197:A:ALA:HA	1:198:A:LEU:HD11	19	3.81
(1,238)	1:9:A:LYS:HG2	1:10:A:LEU:HD11	2	3.79

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PSI	175	50.0	57	32.6	16.3	7	4.0	2.0
PHI	175	50.0	62	35.4	17.7	7	4.0	2.0
Total	350	100.0	119	34.0	34.0	14	4.0	4.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations			Mean (°)	Max (°)	SD (°)	Median (°)
	PSI	PHI	Total				
1	21	26	47	2.18	6.35	1.15	1.76
2	23	28	51	2.3	5.61	0.99	2.1
3	30	22	52	2.31	5.66	1.26	1.9
4	26	23	49	2.15	4.73	1.0	1.82
5	28	21	49	2.1	5.46	1.11	1.66
6	20	22	42	2.11	4.7	0.9	1.93
7	24	28	52	2.41	5.63	1.13	2.14
8	23	27	50	2.27	5.19	0.93	2.16
9	23	29	52	2.26	5.14	1.04	2.04
10	23	28	51	2.27	5.52	1.04	1.88
11	28	22	50	2.11	4.89	0.94	1.9
12	23	22	45	2.3	5.36	1.14	1.82
13	22	23	45	2.37	5.64	1.0	2.25
14	29	26	55	2.2	5.38	1.07	1.85
15	28	24	52	2.18	6.62	1.11	1.92
16	28	25	53	2.34	5.05	0.9	2.23
17	26	29	55	2.05	5.3	1.01	1.69
18	24	29	53	2.17	5.3	0.97	1.77
19	29	26	55	2.15	5.03	1.05	1.78
20	22	28	50	2.3	6.49	1.32	1.9

10.2.1 Bar graph : Dihedral violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints			Fraction of the ensemble	
PSI	PHI	Total	Count ¹	%
9	7	16	1	5.0
5	9	14	2	10.0
7	5	12	3	15.0
0	6	6	4	20.0
4	3	7	5	25.0
2	1	3	6	30.0
4	4	8	7	35.0
2	4	6	8	40.0
0	2	2	9	45.0
1	3	4	10	50.0
1	2	3	11	55.0


Continued on next page...

Continued from previous page...

Number of violated restraints			Fraction of the ensemble	
PSI	PHI	Total	Count ¹	%
2	0	2	12	60.0
4	0	4	13	65.0
2	0	2	14	70.0
2	2	4	15	75.0
1	2	3	16	80.0
1	2	3	17	85.0
1	0	1	18	90.0
2	3	5	19	95.0
7	7	14	20	100.0

¹ Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)




10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

10.4.1 Histogram : Distribution of mean dihedral-angle violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Median
(1,186)	1:101:A:PRO:N	1:101:A:PRO:CA	1:101:A:PRO:C	1:102:A:ARG:N	20	4.71	0.73	4.7
(1,72)	1:42:A:LEU:N	1:42:A:LEU:CA	1:42:A:LEU:C	1:43:A:SER:N	20	4.29	1.37	4.75
(1,102)	1:59:A:GLY:N	1:59:A:GLY:CA	1:59:A:GLY:C	1:60:A:LYS:N	20	4.02	0.73	4.11
(1,129)	1:72:A:VAL:C	1:73:A:PRO:N	1:73:A:PRO:CA	1:73:A:PRO:C	20	3.74	1.08	3.52
(1,233)	1:127:A:PRO:C	1:128:A:GLU:N	1:128:A:GLU:CA	1:128:A:GLU:C	20	3.42	0.61	3.39
(1,176)	1:96:A:LEU:N	1:96:A:LEU:CA	1:96:A:LEU:C	1:97:A:ILE:N	20	3.38	0.61	3.5
(1,22)	1:17:A:PHE:N	1:17:A:PHE:CA	1:17:A:PHE:C	1:18:A:VAL:N	20	3.28	0.39	3.34
(1,231)	1:126:A:GLY:C	1:127:A:PRO:N	1:127:A:PRO:CA	1:127:A:PRO:C	20	2.95	0.67	2.74
(1,219)	1:118:A:THR:C	1:119:A:LEU:N	1:119:A:LEU:CA	1:119:A:LEU:C	20	2.79	0.46	2.86
(1,226)	1:122:A:TYR:N	1:122:A:TYR:CA	1:122:A:TYR:C	1:123:A:VAL:N	20	2.79	0.83	2.63

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,186)	1:101:A:PRO:N	1:101:A:PRO:CA	1:101:A:PRO:C	1:102:A:ARG:N	15	6.62
(1,74)	1:43:A:SER:N	1:43:A:SER:CA	1:43:A:SER:C	1:44:A:THR:N	20	6.49
(1,72)	1:42:A:LEU:N	1:42:A:LEU:CA	1:42:A:LEU:C	1:43:A:SER:N	1	6.35
(1,129)	1:72:A:VAL:C	1:73:A:PRO:N	1:73:A:PRO:CA	1:73:A:PRO:C	20	6.05
(1,129)	1:72:A:VAL:C	1:73:A:PRO:N	1:73:A:PRO:CA	1:73:A:PRO:C	3	5.66
(1,72)	1:42:A:LEU:N	1:42:A:LEU:CA	1:42:A:LEU:C	1:43:A:SER:N	20	5.65
(1,72)	1:42:A:LEU:N	1:42:A:LEU:CA	1:42:A:LEU:C	1:43:A:SER:N	13	5.64
(1,186)	1:101:A:PRO:N	1:101:A:PRO:CA	1:101:A:PRO:C	1:102:A:ARG:N	7	5.63
(1,72)	1:42:A:LEU:N	1:42:A:LEU:CA	1:42:A:LEU:C	1:43:A:SER:N	3	5.62
(1,102)	1:59:A:GLY:N	1:59:A:GLY:CA	1:59:A:GLY:C	1:60:A:LYS:N	2	5.61