



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 4, 2024 – 12:28 PM JST

PDB ID : 7XDJ  
Title : Crystal structure of actinomycin D-echinomycin-d(AGCGCGT/ACGAGCT) complex  
Authors : Chien, C.M.; Satange, R.B.; Hou, M.H.  
Deposited on : 2022-03-27  
Resolution : 2.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	7	<div> <div style="width: 57%; background-color: green;"></div> <div style="width: 43%; background-color: yellow;"></div> </div> <div>57% 43%</div>
1	E	7	<div> <div style="width: 43%; background-color: green;"></div> <div style="width: 57%; background-color: yellow;"></div> </div> <div>43% 57%</div>
1	I	7	<div> <div style="width: 57%; background-color: green;"></div> <div style="width: 43%; background-color: yellow;"></div> </div> <div>57% 43%</div>
1	M	7	<div> <div style="width: 43%; background-color: green;"></div> <div style="width: 57%; background-color: yellow;"></div> </div> <div>43% 57%</div>
2	B	7	<div> <div style="width: 43%; background-color: green;"></div> <div style="width: 57%; background-color: yellow;"></div> </div> <div>43% 57%</div>
2	F	7	<div> <div style="width: 57%; background-color: green;"></div> <div style="width: 29%; background-color: yellow;"></div> <div style="width: 14%; background-color: orange;"></div> </div> <div>57% 29% 14%</div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	J	7	
2	N	7	
3	D	8	
3	H	8	
3	L	8	
3	P	8	
4	C	11	
4	G	11	
4	K	11	
4	O	11	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SAR	C	4	-	-	X	-
4	DVA	G	2	-	-	X	-
4	SAR	G	4	-	-	X	-
4	SAR	K	4	-	-	X	-
4	DVA	O	2	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 1864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P\*AP\*GP\*CP\*GP\*CP\*GP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	7	Total	C	N	O	P	0	0	0
			145	68	28	42	7			
1	E	7	Total	C	N	O	P	0	0	0
			145	68	28	42	7			
1	I	7	Total	C	N	O	P	0	0	0
			145	68	28	42	7			
1	M	7	Total	C	N	O	P	0	0	0
			145	68	28	42	7			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*AP\*CP\*GP\*AP\*GP\*CP\*(BRU))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	7	Total	Br	C	N	O	P	0	0
			144	1	67	28	41	7		
2	F	7	Total	Br	C	N	O	P	0	0
			144	1	67	28	41	7		
2	J	7	Total	Br	C	N	O	P	0	0
			144	1	67	28	41	7		
2	N	7	Total	Br	C	N	O	P	0	0
			144	1	67	28	41	7		

- Molecule 3 is a protein called DSN-ALA-N2C-MVA-DSN-ALA-NCY-MVA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	8	Total	C	N	O	S	0	0	0
			53	33	8	10	2			
3	P	8	Total	C	N	O	S	0	0	0
			53	33	8	10	2			
3	D	8	Total	C	N	O	S	0	0	0
			53	33	8	10	2			
3	H	8	Total	C	N	O	S	0	0	0
			53	33	8	10	2			

- Molecule 4 is a protein called THR-DVA-PRO-SAR-MVA-PXZ-THR-DVA-PRO-SAR-MV A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	11	Total	C	N	O	0	0	0
			90	62	12	16			
4	G	11	Total	C	N	O	0	0	0
			90	62	12	16			
4	K	11	Total	C	N	O	0	0	0
			90	62	12	16			
4	O	11	Total	C	N	O	0	0	0
			90	62	12	16			


- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	J	1	Total	Cl	0	0
			1	1		
5	N	1	Total	Cl	0	0
			1	1		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	J	1	Total	Mg	0	0
			1	1		
6	N	1	Total	Mg	0	0
			1	1		

- Molecule 7 is 2-CARBOXYQUINOXALINE (three-letter code: QUI) (formula: C<sub>9</sub>H<sub>6</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	L	1	Total	C	N	O	0	0
			12	9	2	1		
7	L	1	Total	C	N	O	0	0
			12	9	2	1		
7	P	1	Total	C	N	O	0	0
			12	9	2	1		
7	P	1	Total	C	N	O	0	0
			12	9	2	1		
7	D	1	Total	C	N	O	0	0
			12	9	2	1		
7	D	1	Total	C	N	O	0	0
			12	9	2	1		
7	H	1	Total	C	N	O	0	0
			12	9	2	1		
7	H	1	Total	C	N	O	0	0
			12	9	2	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	4	Total	O	0	0
			4	4		
8	B	4	Total	O	0	0
			4	4		
8	E	3	Total	O	0	0
			3	3		
8	F	4	Total	O	0	0
			4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	I	4	Total 4	O 4	0	0
8	J	1	Total 1	O 1	0	0
8	L	3	Total 3	O 3	0	0
8	M	4	Total 4	O 4	0	0
8	N	1	Total 1	O 1	0	0
8	C	1	Total 1	O 1	0	0
8	G	1	Total 1	O 1	0	0
8	K	1	Total 1	O 1	0	0
8	D	1	Total 1	O 1	0	0
8	H	2	Total 2	O 2	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(P\*AP\*GP\*CP\*GP\*CP\*GP\*T)-3')

Chain A: 



- Molecule 1: DNA (5'-D(P\*AP\*GP\*CP\*GP\*CP\*GP\*T)-3')

Chain E: 



- Molecule 1: DNA (5'-D(P\*AP\*GP\*CP\*GP\*CP\*GP\*T)-3')

Chain I: 



- Molecule 1: DNA (5'-D(P\*AP\*GP\*CP\*GP\*CP\*GP\*T)-3')

Chain M: 



- Molecule 2: DNA (5'-D(P\*AP\*CP\*GP\*AP\*GP\*CP\*(BRU))-3')

Chain B: 




- Molecule 2: DNA (5'-D(P\*AP\*CP\*GP\*AP\*GP\*CP\*(BRU))-3')

Chain F: 






- Molecule 2: DNA (5'-D(P\*AP\*CP\*GP\*AP\*GP\*CP\*(BRU))-3')



- Molecule 2: DNA (5'-D(P\*AP\*CP\*GP\*AP\*GP\*CP\*(BRU))-3')



- Molecule 3: DSN-ALA-N2C-MVA-DSN-ALA-NCY-MVA



- Molecule 3: DSN-ALA-N2C-MVA-DSN-ALA-NCY-MVA




- Molecule 3: DSN-ALA-N2C-MVA-DSN-ALA-NCY-MVA




- Molecule 3: DSN-ALA-N2C-MVA-DSN-ALA-NCY-MVA




- Molecule 4: THR-DVA-PRO-SAR-MVA-PXZ-THR-DVA-PRO-SAR-MVA



- Molecule 4: THR-DVA-PRO-SAR-MVA-PXZ-THR-DVA-PRO-SAR-MVA



- Molecule 4: THR-DVA-PRO-SAR-MVA-PXZ-THR-DVA-PRO-SAR-MVA



- Molecule 4: THR-DVA-PRO-SAR-MVA-PXZ-THR-DVA-PRO-SAR-MVA



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	132.52Å 132.52Å 49.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.84 – 2.44 24.84 – 2.44	Depositor EDS
% Data completeness (in resolution range)	99.5 (24.84-2.44) 93.5 (24.84-2.44)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.44Å)	Xtriage
Refinement program	PHENIX 1.14 _3260	Depositor
R, $R_{free}$	0.201 , 0.236 0.203 , 0.237	Depositor DCC
$R_{free}$ test set	1220 reflections (10.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.0	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 44.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.478 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1864	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DVA, PXZ, N2C, QUI, CL, MVA, DSN, SAR, NCY, MG, BRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.38	0/162	0.92	0/248
1	E	1.37	1/162 (0.6%)	0.99	0/248
1	I	1.56	2/162 (1.2%)	0.82	0/248
1	M	1.02	0/162	1.00	0/248
2	B	1.34	1/139 (0.7%)	0.80	0/212
2	F	1.34	2/139 (1.4%)	0.88	0/212
2	J	1.43	1/139 (0.7%)	0.94	0/212
2	N	0.95	0/139	0.88	0/212
3	D	0.20	0/8	0.26	0/8
3	H	0.46	0/8	0.39	0/8
3	L	0.52	0/8	0.57	0/8
3	P	0.37	0/8	0.59	0/8
4	C	0.51	0/26	1.35	0/30
4	G	0.54	0/26	1.43	0/30
4	K	1.09	0/26	1.33	0/30
4	O	0.66	0/26	1.20	0/30
All	All	1.26	7/1340 (0.5%)	0.93	0/1992

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	C	0	2
4	G	0	2
4	K	0	2
4	O	0	2
All	All	0	8

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	5	DG	O3'-P	-5.89	1.54	1.61
1	E	3	DC	O3'-P	-5.87	1.54	1.61
2	B	3	DG	O3'-P	-5.55	1.54	1.61
1	I	3	DC	O3'-P	-5.42	1.54	1.61
2	F	5	DG	O3'-P	-5.39	1.54	1.61

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	C	2	DVA	Peptide
4	C	8	DVA	Peptide
4	G	2	DVA	Peptide
4	G	8	DVA	Peptide
4	K	2	DVA	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	145	0	79	9	0
1	E	145	0	79	10	0
1	I	145	0	79	5	0
1	M	145	0	79	4	0
2	B	144	0	76	2	0
2	F	144	0	76	1	0
2	J	144	0	75	8	0
2	N	144	0	76	6	0
3	D	53	0	54	1	0
3	H	53	0	54	2	0
3	L	53	0	54	4	0
3	P	53	0	53	1	0
4	C	90	0	84	15	0
4	G	90	0	84	12	0
4	K	90	0	85	10	0
4	O	90	0	85	9	0
5	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	J	1	0	0	0	0
5	N	1	0	0	0	0
6	A	1	0	0	0	0
6	J	1	0	0	0	0
6	N	1	0	0	0	0
7	D	24	0	10	0	0
7	H	24	0	10	1	0
7	L	24	0	10	0	0
7	P	24	0	10	0	0
8	A	4	0	0	0	0
8	B	4	0	0	1	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
8	E	3	0	0	0	0
8	F	4	0	0	0	0
8	G	1	0	0	0	0
8	H	2	0	0	0	0
8	I	4	0	0	0	0
8	J	1	0	0	0	0
8	K	1	0	0	0	0
8	L	3	0	0	0	0
8	M	4	0	0	0	0
8	N	1	0	0	0	0
All	All	1864	0	1212	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:7:DT:H5''	1:E:7:DT:H6	1.30	0.95
4:C:2:DVA:HA	4:C:4:SAR:HA2	1.52	0.91
1:I:4:DG:H21	2:J:4:DA:H62	1.14	0.88
4:G:2:DVA:HA	4:G:4:SAR:HA2	1.56	0.87
2:N:7:BRU:H1'	4:K:4:SAR:HN1	1.58	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	2/8 (25%)	2 (100%)	0	0	100	100
3	H	2/8 (25%)	2 (100%)	0	0	100	100
3	L	2/8 (25%)	2 (100%)	0	0	100	100
3	P	2/8 (25%)	2 (100%)	0	0	100	100
4	C	2/11 (18%)	0	0	2 (100%)	0	0
4	G	2/11 (18%)	0	0	2 (100%)	0	0
4	K	2/11 (18%)	0	0	2 (100%)	0	0
4	O	2/11 (18%)	0	0	2 (100%)	0	0
All	All	16/76 (21%)	8 (50%)	0	8 (50%)	0	0

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	C	3	PRO
4	C	9	PRO
4	G	3	PRO
4	G	9	PRO
4	O	3	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	C	4/4 (100%)	4 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	G	4/4 (100%)	4 (100%)	0	100	100
4	K	4/4 (100%)	4 (100%)	0	100	100
4	O	4/4 (100%)	3 (75%)	1 (25%)	0	0
All	All	16/16 (100%)	15 (94%)	1 (6%)	18	23

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	O	9	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

52 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SAR	C	4	4	4,4,5	1.17	0	1,3,5	3.70	1 (100%)
2	BRU	J	7	2,1	18,21,22	3.39	12 (66%)	26,30,33	3.36	13 (50%)
4	MVA	C	5	4	6,7,8	0.25	0	7,8,10	1.75	1 (14%)
4	SAR	K	4	4	4,4,5	1.47	1 (25%)	1,3,5	3.95	1 (100%)
3	MVA	P	8	3	6,7,8	0.63	0	7,8,10	1.56	2 (28%)
3	N2C	L	3	3	6,7,8	1.24	1 (16%)	5,7,9	1.14	0
3	MVA	H	8	3	6,7,8	0.57	0	7,8,10	2.31	3 (42%)
2	BRU	N	7	2,1	18,21,22	2.77	10 (55%)	26,30,33	3.46	12 (46%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MVA	K	5	4	6,7,8	0.26	0	7,8,10	1.55	1 (14%)
4	SAR	G	4	4	4,4,5	1.21	0	1,3,5	3.51	1 (100%)
3	NCY	D	7	3	5,6,7	1.03	0	5,6,8	1.02	0
2	BRU	F	7	2,1	18,21,22	2.71	10 (55%)	26,30,33	2.98	12 (46%)
4	SAR	G	10	4	4,4,5	0.81	0	1,3,5	1.88	0
4	MVA	G	11	4	6,7,8	0.19	0	7,8,10	1.39	1 (14%)
3	N2C	H	3	3	6,7,8	1.25	1 (16%)	5,7,9	0.84	0
2	BRU	B	7	2,1	18,21,22	3.09	11 (61%)	26,30,33	3.07	12 (46%)
4	MVA	C	11	4	6,7,8	0.29	0	7,8,10	2.02	1 (14%)
3	MVA	L	4	3	6,7,8	0.54	0	7,8,10	1.44	2 (28%)
4	MVA	G	5	4	6,7,8	0.38	0	7,8,10	1.61	2 (28%)
4	SAR	O	4	4	4,4,5	0.94	0	1,3,5	3.53	1 (100%)
3	MVA	P	4	3	6,7,8	0.45	0	7,8,10	1.43	0
3	MVA	H	4	3	6,7,8	0.44	0	7,8,10	1.89	2 (28%)
3	NCY	P	7	3	5,6,7	1.30	1 (20%)	5,6,8	1.36	1 (20%)
3	N2C	P	3	3	6,7,8	1.12	1 (16%)	5,7,9	1.00	0
4	SAR	C	10	4	4,4,5	0.87	0	1,3,5	1.50	0
3	N2C	D	3	3	6,7,8	0.95	0	5,7,9	0.98	0
3	NCY	L	7	3	5,6,7	1.16	1 (20%)	5,6,8	1.33	1 (20%)
3	MVA	D	4	3	6,7,8	0.55	0	7,8,10	1.33	0
4	SAR	K	10	4	4,4,5	0.89	0	1,3,5	1.35	0
4	SAR	O	10	4	4,4,5	0.91	0	1,3,5	1.47	0
3	MVA	D	8	3	6,7,8	0.46	0	7,8,10	1.49	2 (28%)
4	MVA	K	11	4	6,7,8	0.27	0	7,8,10	0.92	0
4	MVA	O	11	4	6,7,8	0.30	0	7,8,10	1.58	1 (14%)
4	MVA	O	5	4	6,7,8	0.37	0	7,8,10	1.30	1 (14%)
3	MVA	L	8	3	6,7,8	0.39	0	7,8,10	1.83	2 (28%)
3	NCY	H	7	3	5,6,7	1.33	1 (20%)	5,6,8	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAR	C	4	4	-	1/1/2/3	-
2	BRU	J	7	2,1	-	1/7/21/22	0/2/2/2
4	MVA	C	5	4	-	1/6/8/10	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAR	K	4	4	-	1/1/2/3	-
3	MVA	P	8	3	-	5/6/8/10	-
3	N2C	L	3	3	-	3/4/7/9	-
3	MVA	H	8	3	-	1/6/8/10	-
2	BRU	N	7	2,1	-	2/7/21/22	0/2/2/2
4	MVA	K	5	4	-	1/6/8/10	-
4	SAR	G	4	4	-	1/1/2/3	-
3	NCY	D	7	3	-	0/3/6/8	-
2	BRU	F	7	2,1	-	1/7/21/22	0/2/2/2
4	SAR	G	10	4	-	1/1/2/3	-
4	MVA	G	11	4	-	2/6/8/10	-
3	N2C	H	3	3	-	3/4/7/9	-
2	BRU	B	7	2,1	-	1/7/21/22	0/2/2/2
4	MVA	C	11	4	-	2/6/8/10	-
3	MVA	L	4	3	-	2/6/8/10	-
4	MVA	G	5	4	-	3/6/8/10	-
4	SAR	O	4	4	-	1/1/2/3	-
3	MVA	P	4	3	-	2/6/8/10	-
3	MVA	H	4	3	-	5/6/8/10	-
3	NCY	P	7	3	-	0/3/6/8	-
3	N2C	P	3	3	-	3/4/7/9	-
4	SAR	C	10	4	-	1/1/2/3	-
3	N2C	D	3	3	-	2/4/7/9	-
3	NCY	L	7	3	-	1/3/6/8	-
3	MVA	D	4	3	-	1/6/8/10	-
4	SAR	K	10	4	-	1/1/2/3	-
4	SAR	O	10	4	-	1/1/2/3	-
3	MVA	D	8	3	-	5/6/8/10	-
4	MVA	K	11	4	-	3/6/8/10	-
4	MVA	O	11	4	-	1/6/8/10	-
4	MVA	O	5	4	-	2/6/8/10	-
3	MVA	L	8	3	-	1/6/8/10	-
3	NCY	H	7	3	-	0/3/6/8	-

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	7	BRU	O4'-C4'	-6.22	1.31	1.45
2	B	7	BRU	C5'-C4'	6.19	1.70	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	7	BRU	C2-N1	-5.94	1.28	1.38
2	N	7	BRU	C2-N1	-5.40	1.29	1.38
2	J	7	BRU	O4'-C1'	-5.04	1.31	1.42

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	7	BRU	O5'-C5'-C4'	9.16	140.14	108.99
2	J	7	BRU	O4'-C4'-C3'	7.03	122.07	105.67
2	B	7	BRU	N3-C2-N1	6.33	123.29	114.89
2	N	7	BRU	O4'-C4'-C5'	6.16	129.63	109.37
2	N	7	BRU	N3-C2-N1	5.87	122.68	114.89

There are no chirality outliers.

5 of 62 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	3	N2C	CA-CB-SG-CD
3	D	3	N2C	CA-CB-SG-CD
3	H	3	N2C	CA-CB-SG-CD
3	L	4	MVA	CB-CA-N-CN
3	P	4	MVA	CB-CA-N-CN

There are no ring outliers.

21 monomers are involved in 39 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	4	SAR	9	0
4	C	5	MVA	1	0
4	K	4	SAR	5	0
3	L	3	N2C	1	0
2	N	7	BRU	2	0
4	K	5	MVA	1	0
4	G	4	SAR	8	0
2	F	7	BRU	1	0
4	G	10	SAR	2	0
4	G	11	MVA	1	0
4	C	11	MVA	2	0
3	L	4	MVA	1	0
4	G	5	MVA	1	0
4	O	4	SAR	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	P	7	NCY	1	0
4	C	10	SAR	2	0
3	D	3	N2C	1	0
4	K	10	SAR	1	0
4	O	10	SAR	1	0
4	O	5	MVA	1	0
3	H	7	NCY	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 6 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
7	QUI	L	101	3	13,13,14	2.19	3 (23%)	17,17,19	2.10	4 (23%)
7	QUI	P	101	3	13,13,14	2.12	3 (23%)	17,17,19	2.29	5 (29%)
7	QUI	P	102	3	13,13,14	2.12	3 (23%)	17,17,19	4.74	5 (29%)
7	QUI	H	101	3	13,13,14	2.31	3 (23%)	17,17,19	2.07	5 (29%)
7	QUI	D	102	3	13,13,14	2.58	3 (23%)	17,17,19	2.03	5 (29%)
7	QUI	L	102	3	13,13,14	2.06	3 (23%)	17,17,19	4.50	6 (35%)
7	QUI	H	102	3	13,13,14	2.09	3 (23%)	17,17,19	2.18	5 (29%)
7	QUI	D	101	-	13,13,14	2.86	3 (23%)	17,17,19	2.05	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	QUI	L	101	3	-	0/2/2/4	0/2/2/2
7	QUI	P	101	3	-	0/2/2/4	0/2/2/2
7	QUI	P	102	3	-	0/2/2/4	0/2/2/2
7	QUI	H	101	3	-	0/2/2/4	0/2/2/2
7	QUI	D	102	3	-	0/2/2/4	0/2/2/2
7	QUI	L	102	3	-	0/2/2/4	0/2/2/2
7	QUI	H	102	3	-	0/2/2/4	0/2/2/2
7	QUI	D	101	-	-	0/2/2/4	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	D	101	QUI	C2-C	7.19	1.56	1.48
7	D	102	QUI	C2-C	6.95	1.55	1.48
7	D	101	QUI	O1-C	6.32	1.43	1.21
7	H	101	QUI	O1-C	6.19	1.42	1.21
7	L	101	QUI	O1-C	5.61	1.40	1.21

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	P	102	QUI	O1-C-C2	-17.80	107.35	124.22
7	L	102	QUI	O1-C-C2	-16.40	108.67	124.22
7	P	101	QUI	C-C2-N1	5.98	120.52	114.66
7	L	101	QUI	C-C2-N1	5.87	120.42	114.66
7	P	102	QUI	C-C2-N1	5.76	120.31	114.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	H	102	QUI	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	7/7 (100%)	-0.84	0 100 100	48, 57, 63, 65	0
1	E	7/7 (100%)	-0.84	0 100 100	48, 58, 67, 76	0
1	I	7/7 (100%)	-0.90	0 100 100	48, 49, 59, 63	0
1	M	7/7 (100%)	-0.85	0 100 100	46, 48, 57, 69	0
2	B	6/7 (85%)	-0.77	0 100 100	53, 55, 64, 77	0
2	F	6/7 (85%)	-0.68	0 100 100	52, 55, 68, 88	0
2	J	6/7 (85%)	-0.74	0 100 100	53, 54, 62, 69	0
2	N	6/7 (85%)	-0.70	0 100 100	52, 53, 55, 75	0
3	D	2/8 (25%)	-0.47	0 100 100	62, 62, 62, 62	0
3	H	2/8 (25%)	-0.34	0 100 100	47, 47, 47, 55	0
3	L	2/8 (25%)	-0.10	0 100 100	47, 47, 47, 48	0
3	P	2/8 (25%)	-0.09	0 100 100	53, 53, 53, 57	0
4	C	4/11 (36%)	0.43	1 (25%) 0 0	47, 47, 56, 80	0
4	G	4/11 (36%)	0.57	1 (25%) 0 0	47, 50, 53, 79	0
4	K	4/11 (36%)	-0.22	0 100 100	48, 57, 60, 78	0
4	O	4/11 (36%)	0.01	0 100 100	50, 54, 58, 82	0
All	All	76/132 (57%)	-0.53	2 (2%) 56 52	46, 55, 78, 88	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	G	3	PRO	3.0
4	C	3	PRO	2.8

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SAR	G	4	5/6	0.86	0.23	69,69,76,78	0
4	SAR	C	4	5/6	0.87	0.20	64,71,75,82	0
4	DVA	K	2	7/8	0.91	0.21	44,61,74,80	0
4	SAR	O	4	5/6	0.91	0.21	62,63,82,90	0
3	NCY	L	7	7/8	0.92	0.10	55,65,67,80	0
4	DVA	G	2	7/8	0.92	0.19	47,51,58,65	0
4	SAR	K	4	5/6	0.92	0.22	53,64,68,71	0
3	DSN	L	1	6/7	0.92	0.14	46,57,64,72	0
4	DVA	C	2	7/8	0.93	0.22	46,48,62,70	0
4	DVA	O	2	7/8	0.93	0.16	45,50,73,74	0
3	DSN	H	5	6/7	0.93	0.12	58,62,66,67	0
4	DVA	K	8	7/8	0.94	0.17	39,49,65,69	0
3	NCY	D	7	7/8	0.95	0.14	62,67,81,84	0
4	MVA	O	5	8/9	0.95	0.20	45,55,62,62	0
3	NCY	P	7	7/8	0.95	0.11	58,66,80,81	0
4	MVA	K	11	8/9	0.95	0.24	44,57,60,60	0
3	N2C	H	3	8/9	0.96	0.13	55,67,80,82	0
3	MVA	P	4	8/9	0.96	0.17	54,63,68,70	0
3	NCY	H	7	7/8	0.96	0.08	56,62,68,76	0
3	MVA	L	8	8/9	0.96	0.20	50,59,69,72	0
3	MVA	P	8	8/9	0.96	0.15	48,56,62,64	0
4	MVA	G	5	8/9	0.96	0.16	45,48,67,73	0
3	DSN	L	5	6/7	0.96	0.11	46,53,55,55	0
4	DVA	C	8	7/8	0.96	0.24	44,56,59,76	0
4	DVA	G	8	7/8	0.96	0.22	50,55,62,69	0
2	BRU	F	7	20/21	0.96	0.12	38,45,67,74	0
4	DVA	O	8	7/8	0.96	0.17	48,54,68,81	0
4	SAR	G	10	5/6	0.96	0.12	38,46,49,59	0
4	SAR	K	10	5/6	0.96	0.17	45,50,63,64	0
3	DSN	P	1	6/7	0.96	0.12	50,56,57,62	0
3	N2C	P	3	8/9	0.97	0.12	51,70,81,81	0
3	N2C	D	3	8/9	0.97	0.13	60,67,76,76	0
2	BRU	B	7	20/21	0.97	0.11	43,48,68,70	0
3	MVA	L	4	8/9	0.97	0.11	43,56,61,62	0
2	BRU	N	7	20/21	0.97	0.11	41,50,67,71	0
3	MVA	H	4	8/9	0.97	0.16	65,67,70,76	0
4	SAR	C	10	5/6	0.97	0.12	49,50,56,64	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	N2C	L	3	8/9	0.97	0.12	55,62,76,85	0
3	MVA	D	8	8/9	0.97	0.16	53,64,68,71	0
4	SAR	O	10	5/6	0.97	0.13	44,45,48,52	0
4	MVA	C	11	8/9	0.97	0.17	39,46,54,55	0
4	MVA	G	11	8/9	0.97	0.17	40,48,57,60	0
4	MVA	C	5	8/9	0.97	0.14	47,50,60,63	0
2	BRU	J	7	20/21	0.98	0.11	43,54,71,72	0
3	DSN	D	1	6/7	0.98	0.11	50,56,59,72	0
3	DSN	P	5	6/7	0.98	0.10	53,60,69,70	0
4	MVA	K	5	8/9	0.98	0.17	36,44,51,55	0
3	DSN	D	5	6/7	0.98	0.10	67,69,73,74	0
3	DSN	H	1	6/7	0.98	0.11	46,49,56,57	0
3	MVA	D	4	8/9	0.98	0.18	62,68,73,76	0
3	MVA	H	8	8/9	0.98	0.14	42,52,62,70	0
4	MVA	O	11	8/9	0.98	0.16	42,48,54,56	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	MG	N	102	1/1	0.90	0.38	49,49,49,49	0
6	MG	A	102	1/1	0.93	0.20	42,42,42,42	0
6	MG	J	102	1/1	0.94	0.45	53,53,53,53	0
5	CL	A	101	1/1	0.94	0.21	88,88,88,88	0
7	QUI	P	102	12/13	0.95	0.14	43,52,61,62	0
7	QUI	D	101	12/13	0.96	0.20	54,58,65,68	0
5	CL	N	101	1/1	0.97	0.12	62,62,62,62	0
7	QUI	L	101	12/13	0.97	0.13	39,42,54,56	0
7	QUI	D	102	12/13	0.97	0.14	64,70,72,73	0
7	QUI	H	101	12/13	0.97	0.15	41,44,48,53	0
7	QUI	L	102	12/13	0.98	0.14	41,42,45,46	0
7	QUI	P	101	12/13	0.98	0.15	51,52,58,59	0
7	QUI	H	102	12/13	0.98	0.14	48,50,54,54	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CL	J	101	1/1	0.99	0.14	59,59,59,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.