



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2024 – 11:43 AM EDT

PDB ID : 4YCS
Title : Crystal structure of putative lipoprotein from *Peptoclostridium difficile* 630 (fragment)
Authors : Michalska, K.; Wu, R.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2015-02-20
Resolution : 1.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1356 (1.98-1.98)
Clashscore	180529	1437 (1.98-1.98)
Ramachandran outliers	177936	1426 (1.98-1.98)
Sidechain outliers	177891	1426 (1.98-1.98)
RSRZ outliers	164620	1356 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	 2% 93% 5% 2%
1	B	128	 2% 93% 5% 2%
1	C	128	 2% 91% 5% 2%
1	D	128	 3% 89% 6% 5%
1	E	128	 2% 88% 7% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	128	<div><div></div><div>2%</div><div>92%</div><div>5%</div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	123	Total	C	N	O	S	Se	0	0	0
			950	593	155	197	1	4			
1	B	123	Total	C	N	O	S	Se	0	0	0
			950	593	155	197	1	4			
1	C	122	Total	C	N	O	S	Se	0	1	0
			952	594	156	197	1	4			
1	D	122	Total	C	N	O	S	Se	0	4	0
			974	606	159	203	1	5			
1	E	122	Total	C	N	O	S	Se	0	1	0
			953	595	156	197	1	4			
1	F	122	Total	C	N	O	S	Se	0	1	0
			952	595	155	196	1	5			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	42	SER	-	expression tag	UNP A0A031WBX8
A	43	ASN	-	expression tag	UNP A0A031WBX8
A	44	ALA	-	expression tag	UNP A0A031WBX8
B	42	SER	-	expression tag	UNP A0A031WBX8
B	43	ASN	-	expression tag	UNP A0A031WBX8
B	44	ALA	-	expression tag	UNP A0A031WBX8
C	42	SER	-	expression tag	UNP A0A031WBX8
C	43	ASN	-	expression tag	UNP A0A031WBX8
C	44	ALA	-	expression tag	UNP A0A031WBX8
D	42	SER	-	expression tag	UNP A0A031WBX8
D	43	ASN	-	expression tag	UNP A0A031WBX8
D	44	ALA	-	expression tag	UNP A0A031WBX8
E	42	SER	-	expression tag	UNP A0A031WBX8
E	43	ASN	-	expression tag	UNP A0A031WBX8
E	44	ALA	-	expression tag	UNP A0A031WBX8
F	42	SER	-	expression tag	UNP A0A031WBX8
F	43	ASN	-	expression tag	UNP A0A031WBX8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	44	ALA	-	expression tag	UNP A0A031WBX8

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	38	Total	O	0	0
			38	38		
5	B	62	Total	O	0	0
			62	62		
5	C	66	Total	O	0	1
			67	67		
5	D	54	Total	O	0	0
			54	54		
5	E	74	Total	O	0	2
			74	74		
5	F	59	Total	O	0	0
			59	59		

- Molecule 1: Uncharacterized protein



Sequence logo for the 10th position. The y-axis represents information content in bits, ranging from 0 to 0.4. The x-axis shows the amino acid sequence: SER, ASN, ALA, SER, ASN, S47, L80, L121, F133, Y152, S157, A165, T169. The 'S' (Serine) at position 10 has the highest information content, around 0.35 bits. Other residues like 'Y' (Tyrosine) and 'T' (Threonine) also show significant information content.

Sequence logo for the 10th position. The y-axis represents information content in bits, ranging from 0 to 1. The x-axis shows the amino acid sequence: SER, ASN, ALA, SER, ASN, SER, K48, N61, L80, K86, V113, I136, Y152, A165, T169. The 'Y152' position has a red dot above it, indicating a high information content.

Dataset	Nodes
SER	100
ASN	100
ALA	100
SER	100
ASN	100
SER	100
K48	100
N74	100
Q77	100
L80	100
D83	100
T84	100
N85	100
Y152	100
E153	100
D154	100
N155	100
K156	100
S157	100
G158	100
Y159	100
A165	100
T169	100

SER
ASN
ALA
SER
ASN
SER
K48
Q77
D83
T84
N85
K86
V113
I136
I150
S151
Y152
S157
L163
I164
M168
T169



WORLD WIDE
PDB
PROTEIN DATA BANK



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	37.76Å 119.50Å 185.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.98 30.00 – 1.98	Depositor EDS
% Data completeness (in resolution range)	95.9 (30.00-1.98) 95.9 (30.00-1.98)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.180 , 0.226 0.188 , 0.236	Depositor DCC
R_{free} test set	1080 reflections (1.88%)	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6108	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/957	0.66	0/1282
1	B	0.69	0/957	0.70	0/1282
1	C	0.73	0/959	0.70	0/1285
1	D	0.74	0/981	0.69	0/1314
1	E	0.82	0/960	0.75	0/1286
1	F	0.71	0/959	0.70	0/1284
All	All	0.73	0/5773	0.70	0/7733

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	945	2	0
1	B	950	0	945	2	0
1	C	952	0	945	2	0
1	D	974	0	961	6	0
1	E	953	0	947	6	0
1	F	952	0	948	2	0
2	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	6	0	8	0	0
2	D	6	0	7	0	0
3	D	4	0	3	0	0
4	D	1	0	0	0	0
5	A	38	0	0	0	0
5	B	62	0	0	0	0
5	C	67	0	0	0	0
5	D	54	0	0	0	0
5	E	74	0	0	0	0
5	F	59	0	0	0	0
All	All	6108	0	5717	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:VAL:HG22	1:A:136:ILE:HD12	1.88	0.56
1:D:80:LEU:HD11	1:D:165:ALA:HA	1.88	0.55
1:A:150:ILE:HD11	1:A:163:LEU:HD22	1.89	0.54
1:B:80:LEU:HD11	1:B:165:ALA:HA	1.90	0.53
1:C:80:LEU:HD21	1:C:165:ALA:HA	1.93	0.50

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/128 (94%)	120 (99%)	1 (1%)	0	100	100
1	B	121/128 (94%)	120 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	121/128 (94%)	121 (100%)	0	0	100	100
1	D	124/128 (97%)	122 (98%)	2 (2%)	0	100	100
1	E	121/128 (94%)	120 (99%)	1 (1%)	0	100	100
1	F	121/128 (94%)	121 (100%)	0	0	100	100
All	All	729/768 (95%)	724 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/107 (100%)	107 (100%)	0	100	100
1	B	107/107 (100%)	107 (100%)	0	100	100
1	C	107/107 (100%)	104 (97%)	3 (3%)	38	28
1	D	110/107 (103%)	110 (100%)	0	100	100
1	E	107/107 (100%)	104 (97%)	3 (3%)	38	28
1	F	107/107 (100%)	107 (100%)	0	100	100
All	All	645/642 (100%)	639 (99%)	6 (1%)	79	75

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	77[A]	GLN
1	E	77[B]	GLN
1	E	86	LYS
1	C	80	LEU
1	C	61	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	D	201	-	3,3,3	0.77	0	3,3,3	0.89	0
2	GOL	A	201	-	5,5,5	0.29	0	5,5,5	0.40	0
2	GOL	D	203	4	5,5,5	0.46	0	5,5,5	0.54	0
2	GOL	C	201	-	5,5,5	0.33	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	201	-	-	2/4/4/4	-
2	GOL	D	203	4	-	1/4/4/4	-
2	GOL	C	201	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	GOL	C1-C2-C3-O3
2	C	201	GOL	O1-C1-C2-C3
2	A	201	GOL	O2-C2-C3-O3
2	C	201	GOL	O1-C1-C2-O2
2	D	203	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	119/128 (92%)	0.33	2 (1%) 69 77	24, 45, 75, 87	0
1	B	119/128 (92%)	0.12	3 (2%) 58 68	23, 38, 66, 79	0
1	C	118/128 (92%)	-0.02	2 (1%) 69 77	15, 34, 64, 68	1 (0%)
1	D	118/128 (92%)	0.02	4 (3%) 48 58	18, 34, 66, 88	3 (2%)
1	E	118/128 (92%)	-0.07	3 (2%) 58 68	17, 32, 59, 81	1 (0%)
1	F	118/128 (92%)	0.01	2 (1%) 69 77	23, 36, 58, 65	0
All	All	710/768 (92%)	0.07	16 (2%) 61 70	15, 37, 66, 88	5 (0%)

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	THR	4.1
1	D	152	TYR	3.4
1	A	47	SER	3.1
1	B	169	THR	3.0
1	C	169	THR	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	201	6/6	0.77	0.12	75,80,81,82	0
3	ACT	D	201	4/4	0.84	0.20	57,62,66,67	0
2	GOL	D	203	6/6	0.88	0.12	52,64,68,73	0
2	GOL	C	201	6/6	0.88	0.20	58,62,65,66	0
4	NA	D	202	1/1	0.94	0.10	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.