



wwPDB EM Validation Summary Report ⓘ

Jan 1, 2025 – 06:53 PM EST

PDB ID : 8YPK
EMDB ID : EMD-39482
Title : mouse proteasome 20S subunit in complex with compound 1
Authors : Kashima, A.; Arai, Y.
Deposited on : 2024-03-17
Resolution : 2.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY










The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	
1	F	205	
2	P	234	
2	b	234	
3	I	248	
3	N	248	
4	K	246	
4	R	246	
5	T	201	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	V	201	
6	S	240	
6	X	240	
7	C	205	
7	D	205	
8	J	255	
8	Q	255	
9	G	263	
9	L	263	
10	H	241	
10	M	241	
11	O	261	
11	Z	261	
12	B	234	
12	E	234	
13	W	264	
13	a	264	
14	U	205	
14	Y	205	

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 47992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	202	Total	C	N	O	S	0	0
			1519	952	259	296	12		
1	F	202	Total	C	N	O	S	0	0
			1519	952	259	296	12		

- Molecule 2 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	P	227	Total	C	N	O	S	0	0
			1782	1138	304	334	6		
2	b	227	Total	C	N	O	S	0	0
			1782	1138	304	334	6		

- Molecule 3 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	N	232	Total	C	N	O	S	0	0
			1819	1146	322	346	5		
3	I	232	Total	C	N	O	S	0	0
			1819	1146	322	346	5		

- Molecule 4 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	R	236	Total	C	N	O	S	0	0
			1831	1163	304	351	13		
4	K	236	Total	C	N	O	S	0	0
			1831	1163	304	351	13		

- Molecule 5 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	T	196	Total	C	N	O	S	0	0
			1570	1006	267	288	9		
5	V	196	Total	C	N	O	S	0	0
			1570	1006	267	288	9		

- Molecule 6 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	S	212	Total	C	N	O	S	0	0
			1643	1041	280	312	10		
6	X	212	Total	C	N	O	S	0	0
			1643	1041	280	312	10		

- Molecule 7 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	D	200	Total	C	N	O	S	0	0
			1551	980	270	292	9		
7	C	200	Total	C	N	O	S	0	0
			1551	980	270	292	9		

- Molecule 8 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	J	235	Total	C	N	O	S	0	0
			1840	1168	315	346	11		
8	Q	235	Total	C	N	O	S	0	0
			1840	1168	315	346	11		

- Molecule 9 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	L	234	Total	C	N	O	S	0	0
			1836	1150	328	347	11		
9	G	234	Total	C	N	O	S	0	0
			1836	1150	328	347	11		

- Molecule 10 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	H	231	Total	C	N	O	S	0	0
			1761	1106	292	352	11		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
10	M	231	Total	C	N	O	S	0	0
			1761	1106	292	352	11		

- Molecule 11 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	O	239	Total	C	N	O	S	0	0
			1881	1191	321	359	10		
11	Z	239	Total	C	N	O	S	0	0
			1881	1191	321	359	10		

- Molecule 12 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	B	220	Total	C	N	O	S	0	0
			1656	1044	282	318	12		
12	E	220	Total	C	N	O	S	0	0
			1656	1044	282	318	12		


- Molecule 13 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	W	214	Total	C	N	O	S	0	0
			1671	1054	289	316	12		
13	a	214	Total	C	N	O	S	0	0
			1671	1054	289	316	12		

- Molecule 14 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	Y	204	Total	C	N	O	S	0	0
			1592	1013	265	295	19		
14	U	204	Total	C	N	O	S	0	0
			1592	1013	265	295	19		

- Molecule 15 is [(1R)-1-[(1-cyclohexyl-1,2,3-triazol-4-yl)carbonylamino]-3-methyl-butyl]boronic acid (three-letter code: A1L0C) (formula: C₁₄H₂₅BN₄O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
15	A	1	Total	B	C	N	O	0
			22	1	14	4	3	
15	D	1	Total	B	C	N	O	0
			22	1	14	4	3	
15	C	1	Total	B	C	N	O	0
			22	1	14	4	3	
15	F	1	Total	B	C	N	O	0
			22	1	14	4	3	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit beta type-6




- Molecule 1: Proteasome subunit beta type-6




- Molecule 2: Proteasome subunit alpha type-2





- Molecule 2: Proteasome subunit alpha type-2









• Molecule 3: Proteasome subunit alpha type-7




• Molecule 3: Proteasome subunit alpha type-7



• Molecule 4: Proteasome subunit alpha type-6





• Molecule 4: Proteasome subunit alpha type-6








- Molecule 5: Proteasome subunit beta type-2





- Molecule 5: Proteasome subunit beta type-2




- Molecule 6: Proteasome subunit beta type-1




- Molecule 6: Proteasome subunit beta type-1



- Molecule 7: Proteasome subunit beta type-5





- Molecule 7: Proteasome subunit beta type-5








- Molecule 8: Proteasome subunit alpha type-3




- Molecule 8: Proteasome subunit alpha type-3




- Molecule 9: Proteasome subunit alpha type-1




- Molecule 9: Proteasome subunit alpha type-1





- Molecule 10: Proteasome subunit alpha type-5

Chain H: 




• Molecule 10: Proteasome subunit alpha type-5

Chain M: 




• Molecule 11: Proteasome subunit alpha type-4

Chain O: 





• Molecule 11: Proteasome subunit alpha type-4

Chain Z: 





• Molecule 12: Proteasome subunit beta type-7

Chain B: 




- Molecule 12: Proteasome subunit beta type-7

Chain E: 




- Molecule 13: Proteasome subunit beta type-4

Chain W: 





- Molecule 13: Proteasome subunit beta type-4

Chain a: 





- Molecule 14: Proteasome subunit beta type-3

Chain Y: 



● Molecule 14: Proteasome subunit beta type-3

Chain U:  86% 14%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	25513	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	49.692	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.211	Depositor
Minimum map value	-0.122	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.0343	Depositor
Map size (Å)	292.16095, 292.16095, 292.16095	wwPDB
Map dimensions	254, 254, 254	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.15024, 1.15024, 1.15024	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1L0C

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/1546	0.46	0/2094
1	F	0.28	0/1546	0.46	0/2094
2	P	0.25	0/1821	0.43	0/2466
2	b	0.26	0/1821	0.43	0/2466
3	I	0.25	0/1846	0.44	0/2495
3	N	0.25	0/1846	0.44	0/2495
4	K	0.25	0/1863	0.44	0/2518
4	R	0.25	0/1863	0.44	0/2518
5	T	0.27	0/1602	0.45	0/2167
5	V	0.27	0/1602	0.45	0/2167
6	S	0.27	0/1674	0.45	0/2257
6	X	0.27	0/1674	0.45	0/2257
7	C	0.28	0/1582	0.41	0/2137
7	D	0.28	0/1582	0.41	0/2137
8	J	0.25	0/1875	0.41	0/2524
8	Q	0.25	0/1875	0.41	0/2524
9	G	0.25	0/1870	0.42	0/2528
9	L	0.25	0/1870	0.42	0/2528
10	H	0.24	0/1788	0.42	0/2415
10	M	0.25	0/1788	0.42	0/2415
11	O	0.27	0/1910	0.49	0/2573
11	Z	0.27	0/1910	0.49	0/2573
12	B	0.26	0/1683	0.45	0/2276
12	E	0.26	0/1683	0.45	0/2276
13	W	0.27	0/1704	0.45	0/2306
13	a	0.27	0/1704	0.45	0/2306
14	U	0.27	0/1621	0.46	0/2185
14	Y	0.27	0/1621	0.46	0/2185
All	All	0.26	0/48770	0.44	0/65882

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/205 (98%)	188 (94%)	11 (6%)	1 (0%)	25	49
1	F	200/205 (98%)	188 (94%)	11 (6%)	1 (0%)	25	49
2	P	225/234 (96%)	206 (92%)	16 (7%)	3 (1%)	10	26
2	b	225/234 (96%)	206 (92%)	16 (7%)	3 (1%)	10	26
3	I	230/248 (93%)	217 (94%)	13 (6%)	0	100	100
3	N	230/248 (93%)	217 (94%)	13 (6%)	0	100	100
4	K	232/246 (94%)	215 (93%)	16 (7%)	1 (0%)	30	55
4	R	232/246 (94%)	215 (93%)	16 (7%)	1 (0%)	30	55
5	T	194/201 (96%)	184 (95%)	10 (5%)	0	100	100
5	V	194/201 (96%)	184 (95%)	10 (5%)	0	100	100
6	S	210/240 (88%)	193 (92%)	15 (7%)	2 (1%)	13	33
6	X	210/240 (88%)	193 (92%)	15 (7%)	2 (1%)	13	33
7	C	198/205 (97%)	185 (93%)	13 (7%)	0	100	100
7	D	198/205 (97%)	185 (93%)	13 (7%)	0	100	100
8	J	233/255 (91%)	217 (93%)	16 (7%)	0	100	100
8	Q	233/255 (91%)	218 (94%)	15 (6%)	0	100	100
9	G	232/263 (88%)	215 (93%)	16 (7%)	1 (0%)	30	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
9	L	232/263 (88%)	215 (93%)	16 (7%)	1 (0%)	30	55
10	H	229/241 (95%)	215 (94%)	14 (6%)	0	100	100
10	M	229/241 (95%)	215 (94%)	14 (6%)	0	100	100
11	O	235/261 (90%)	224 (95%)	10 (4%)	1 (0%)	30	55
11	Z	235/261 (90%)	224 (95%)	10 (4%)	1 (0%)	30	55
12	B	218/234 (93%)	206 (94%)	11 (5%)	1 (0%)	25	49
12	E	218/234 (93%)	206 (94%)	11 (5%)	1 (0%)	25	49
13	W	212/264 (80%)	195 (92%)	16 (8%)	1 (0%)	25	49
13	a	212/264 (80%)	195 (92%)	16 (8%)	1 (0%)	25	49
14	U	202/205 (98%)	180 (89%)	20 (10%)	2 (1%)	13	33
14	Y	202/205 (98%)	180 (89%)	20 (10%)	2 (1%)	13	33
All	All	6100/6604 (92%)	5681 (93%)	393 (6%)	26 (0%)	32	55

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	P	40	ALA
2	b	40	ALA
11	O	59	VAL
11	Z	59	VAL
14	Y	78	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/162 (98%)	128 (80%)	31 (20%)	1	3
1	F	159/162 (98%)	128 (80%)	31 (20%)	1	3
2	P	188/191 (98%)	143 (76%)	45 (24%)	0	2
2	b	188/191 (98%)	143 (76%)	45 (24%)	0	2
3	I	195/211 (92%)	151 (77%)	44 (23%)	1	2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	N	195/211 (92%)	151 (77%)	44 (23%)	1	2
4	K	201/210 (96%)	164 (82%)	37 (18%)	1	3
4	R	201/210 (96%)	164 (82%)	37 (18%)	1	3
5	T	166/171 (97%)	147 (89%)	19 (11%)	4	11
5	V	166/171 (97%)	147 (89%)	19 (11%)	4	11
6	S	177/198 (89%)	144 (81%)	33 (19%)	1	3
6	X	177/198 (89%)	144 (81%)	33 (19%)	1	3
7	C	156/161 (97%)	131 (84%)	25 (16%)	2	5
7	D	156/161 (97%)	131 (84%)	25 (16%)	2	5
8	J	192/211 (91%)	155 (81%)	37 (19%)	1	3
8	Q	192/211 (91%)	155 (81%)	37 (19%)	1	3
9	G	200/224 (89%)	155 (78%)	45 (22%)	1	2
9	L	200/224 (89%)	155 (78%)	45 (22%)	1	2
10	H	193/203 (95%)	144 (75%)	49 (25%)	0	1
10	M	193/203 (95%)	144 (75%)	49 (25%)	0	1
11	O	201/221 (91%)	151 (75%)	50 (25%)	0	1
11	Z	201/221 (91%)	151 (75%)	50 (25%)	0	1
12	B	181/195 (93%)	141 (78%)	40 (22%)	1	2
12	E	181/195 (93%)	142 (78%)	39 (22%)	1	2
13	W	176/212 (83%)	142 (81%)	34 (19%)	1	3
13	a	176/212 (83%)	142 (81%)	34 (19%)	1	3
14	U	174/175 (99%)	148 (85%)	26 (15%)	2	6
14	Y	174/175 (99%)	149 (86%)	25 (14%)	2	7
All	All	5118/5490 (93%)	4090 (80%)	1028 (20%)	3	3

5 of 1028 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	Z	180	LYS
14	Y	80	GLN
11	Z	176	LYS
9	L	101	ARG
9	L	38	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 96 such sidechains are listed below:

Mol	Chain	Res	Type
12	B	40	ASN
8	Q	22	GLN
10	M	227	HIS
9	G	5	GLN
13	a	147	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
15	A1L0C	F	301	1	16,23,23	0.63	1 (6%)	20,31,31	1.52	1 (5%)
15	A1L0C	C	301	7	16,23,23	0.61	0	20,31,31	1.59	1 (5%)
15	A1L0C	A	301	1	16,23,23	0.62	1 (6%)	20,31,31	1.53	1 (5%)
15	A1L0C	D	301	7	16,23,23	0.61	0	20,31,31	1.58	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	A1L0C	F	301	1	-	2/9/28/28	0/2/2/2
15	A1L0C	C	301	7	-	1/9/28/28	0/2/2/2
15	A1L0C	A	301	1	-	2/9/28/28	0/2/2/2
15	A1L0C	D	301	7	-	1/9/28/28	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	F	301	A1L0C	N3-N2	-2.09	1.30	1.34
15	A	301	A1L0C	N3-N2	-2.07	1.30	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	C	301	A1L0C	C2-N1-C9	6.23	131.00	125.48
15	D	301	A1L0C	C2-N1-C9	6.18	130.96	125.48
15	A	301	A1L0C	C2-N1-C9	5.99	130.79	125.48
15	F	301	A1L0C	C2-N1-C9	5.98	130.78	125.48

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
15	A	301	A1L0C	O3-C3-N4-C4
15	D	301	A1L0C	C10-C9-N1-N2
15	C	301	A1L0C	C10-C9-N1-N2
15	F	301	A1L0C	O3-C3-N4-C4
15	A	301	A1L0C	C1-C3-N4-C4


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-39482. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 127




Y Index: 127




Z Index: 127


6.2.2 Raw map



X Index: 127



Y Index: 127




Z Index: 127


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 143




Y Index: 110




Z Index: 146


6.3.2 Raw map



X Index: 143



Y Index: 110




Z Index: 146


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0343. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_39482_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 254 nm³; this corresponds to an approximate mass of 229 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.370 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.370 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.70	-	-
Author-provided FSC curve	2.72	3.29	2.81
Unmasked-calculated*	3.25	4.04	3.31

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.25 differs from the reported value 2.7 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-39482 and PDB model 8YPK. Per-residue inclusion information can be found in [section 3](#) on [page 8](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0343 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0343).



























































9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0343) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8450	 0.6210
A	 0.8910	 0.6400
B	 0.8720	 0.6390
C	 0.9050	 0.6460
D	 0.9020	 0.6460
E	 0.8710	 0.6390
F	 0.8910	 0.6400
G	 0.8200	 0.6090
H	 0.8050	 0.6050
I	 0.8080	 0.5970
J	 0.7800	 0.5980
K	 0.7760	 0.6020
L	 0.8210	 0.6110
M	 0.8070	 0.6050
N	 0.8100	 0.5990
O	 0.8000	 0.5930
P	 0.8220	 0.6100
Q	 0.7840	 0.5990
R	 0.7730	 0.6010
S	 0.8770	 0.6350
T	 0.9100	 0.6490
U	 0.9080	 0.6500
V	 0.9090	 0.6480
W	 0.9010	 0.6440
X	 0.8750	 0.6340
Y	 0.9090	 0.6490
Z	 0.8000	 0.5910
a	 0.8980	 0.6440
b	 0.8240	 0.6100

