



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2025 – 12:44 PM JST

PDB ID : 8YU4
Title : Structure of cyclohexanone monooxygenase mutant from *Acinetobacter calcoaceticus*
Authors : Qiang, G.; Zheng, Y.C.; Feng, L.; Yu, H.L.
Deposited on : 2024-03-26
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.21
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	12358 (2.04-2.00)
Clashscore	180529	13897 (2.04-2.00)
Ramachandran outliers	177936	13770 (2.04-2.00)
Sidechain outliers	177891	13769 (2.04-2.00)
RSRZ outliers	164620	12358 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	548	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative flavin-binding monooxygenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	520	Total	C	N	O	S	0	0	0
			4143	2639	692	794	18			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	initiating methionine	UNP A0A0A8XFY0
A	-4	HIS	-	expression tag	UNP A0A0A8XFY0
A	-3	HIS	-	expression tag	UNP A0A0A8XFY0
A	-2	HIS	-	expression tag	UNP A0A0A8XFY0
A	-1	HIS	-	expression tag	UNP A0A0A8XFY0
A	0	HIS	-	expression tag	UNP A0A0A8XFY0
A	1	HIS	-	expression tag	UNP A0A0A8XFY0
A	143	PRO	LEU	engineered mutation	UNP A0A0A8XFY0
A	145	SER	ALA	engineered mutation	UNP A0A0A8XFY0
A	146	SER	ALA	engineered mutation	UNP A0A0A8XFY0
A	149	TRP	LEU	engineered mutation	UNP A0A0A8XFY0
A	151	ILE	LYS	engineered mutation	UNP A0A0A8XFY0
A	246	TYR	PHE	engineered mutation	UNP A0A0A8XFY0
A	326	CYS	LYS	engineered mutation	UNP A0A0A8XFY0
A	386	SER	ASN	engineered mutation	UNP A0A0A8XFY0
A	388	LYS	ILE	engineered mutation	UNP A0A0A8XFY0
A	390	ILE	MET	engineered mutation	UNP A0A0A8XFY0
A	426	PHE	LEU	engineered mutation	UNP A0A0A8XFY0
A	432	LEU	PHE	engineered mutation	UNP A0A0A8XFY0
A	433	ALA	THR	engineered mutation	UNP A0A0A8XFY0
A	435	SER	LEU	engineered mutation	UNP A0A0A8XFY0
A	438	ILE	SER	engineered mutation	UNP A0A0A8XFY0
A	488	LYS	GLU	engineered mutation	UNP A0A0A8XFY0
A	489	CYS	SER	engineered mutation	UNP A0A0A8XFY0
A	490	ARG	TRP	engineered mutation	UNP A0A0A8XFY0
A	505	LEU	PHE	engineered mutation	UNP A0A0A8XFY0

- # NAI

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $\text{C}_{27}\text{H}_{33}\text{N}_9\text{O}_{15}\text{P}_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	216	Total 216	O 216	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative flavin-binding monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.92Å 53.23Å 101.49Å 90.00° 96.86° 90.00°	Depositor
Resolution (Å)	44.46 – 2.02 44.46 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.46-2.02) 92.2 (44.46-2.02)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.01Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.218 , 0.260 0.219 , 0.259	Depositor DCC
R_{free} test set	1754 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4456	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, NAI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/4237	0.45	0/5735

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4143	0	4043	31	0
2	A	44	0	27	1	0
3	A	53	0	31	1	0
4	A	216	0	0	0	0
All	All	4456	0	4101	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:SER:HB3	1:A:381:ASP:HB2	1.73	0.69
1:A:326:CYS:SG	1:A:327:ARG:N	2.68	0.65
1:A:41:GLU:HG2	1:A:92:LYS:HE2	1.79	0.63
1:A:395:LYS:O	1:A:397:GLY:N	2.34	0.61
1:A:155:ILE:HA	1:A:356:ILE:HD11	1.86	0.57
1:A:395:LYS:O	1:A:398:ILE:HD12	2.06	0.56
1:A:174:PHE:HB3	1:A:199:VAL:HG12	1.88	0.55
1:A:391:ASP:OD1	1:A:393:ARG:NH1	2.39	0.55
1:A:395:LYS:C	1:A:397:GLY:H	2.13	0.53
1:A:348:VAL:HG22	1:A:352:PRO:HA	1.91	0.52
1:A:11:ILE:HG12	1:A:138:ILE:HB	1.93	0.50
1:A:119:ASN:N	1:A:119:ASN:OD1	2.46	0.48
1:A:71:LEU:HD21	1:A:94:VAL:HG22	1.96	0.47
1:A:49:ASN:O	1:A:84:GLN:HG3	2.14	0.47
1:A:298:GLN:HE22	1:A:324:TYR:HA	1.79	0.47
1:A:476:SER:HB3	1:A:503:VAL:HG21	1.97	0.47
1:A:294:ASN:OD1	1:A:298:GLN:NE2	2.41	0.47
1:A:57:ASP:HB2	2:A:601:NAI:O7N	2.15	0.47
1:A:227:LYS:HG3	1:A:228:ILE:N	2.29	0.46
1:A:149:TRP:CZ2	1:A:164:HIS:HB2	2.52	0.45
1:A:482:THR:HG22	1:A:484:PHE:H	1.82	0.45
1:A:167:ARG:HA	1:A:167:ARG:HD2	1.72	0.44
3:A:602:FAD:H9	3:A:602:FAD:H1'1	1.77	0.44
1:A:16:PHE:HB3	1:A:439:ILE:HD13	2.00	0.43
1:A:317:LYS:HD3	1:A:347:ASP:HB2	2.01	0.43
1:A:486:LYS:HB2	1:A:486:LYS:HE2	1.76	0.43
1:A:88:LEU:HG	1:A:92:LYS:HE3	2.01	0.42
1:A:384:ASP:OD1	1:A:384:ASP:N	2.54	0.41
1:A:460:GLN:HB3	1:A:531:LYS:HB2	2.01	0.41
1:A:120:SER:OG	1:A:460:GLN:OE1	2.39	0.41
1:A:298:GLN:NE2	1:A:324:TYR:HA	2.35	0.40
1:A:96:ASP:OD1	1:A:101:ARG:NH1	2.41	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/548 (94%)	500 (97%)	13 (2%)	3 (1%)	22	16

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	396	ASP
1	A	5	MET
1	A	326	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	449/474 (95%)	431 (96%)	18 (4%)	27	24

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	4	LYS
1	A	34	ARG
1	A	119	ASN
1	A	120	SER
1	A	121	PHE
1	A	128	ASN
1	A	167	ARG
1	A	208	SER
1	A	223	THR
1	A	227	LYS
1	A	232	TYR
1	A	309	VAL
1	A	348	VAL
1	A	396	ASP
1	A	454	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	456	HIS
1	A	477	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	A	602	-	53,58,58	1.29	5 (9%)	68,89,89	1.34	10 (14%)
2	NAI	A	601	-	42,48,48	0.99	2 (4%)	47,73,73	1.24	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	602	-	-	16/30/50/50	0/6/6/6
2	NAI	A	601	-	-	9/25/72/72	0/5/5/5

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	FAD	C9A-C5X	5.08	1.49	1.41
2	A	601	NAI	C6N-C5N	3.41	1.39	1.33
3	A	602	FAD	C8-C7	3.30	1.49	1.40
3	A	602	FAD	C4-N3	-2.55	1.34	1.38
3	A	602	FAD	C4X-N5	2.44	1.35	1.30
2	A	601	NAI	C5A-C4A	2.44	1.47	1.40
3	A	602	FAD	C5A-C4A	2.41	1.47	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	FAD	N3A-C2A-N1A	-3.33	123.48	128.68
2	A	601	NAI	N3A-C2A-N1A	-3.28	123.55	128.68
2	A	601	NAI	PN-O3-PA	-3.00	122.52	132.83
3	A	602	FAD	C4-C4X-N5	2.90	122.36	118.23
3	A	602	FAD	C4X-C10-N1	-2.79	118.25	124.73
3	A	602	FAD	C4A-C5A-N7A	-2.75	106.53	109.40
3	A	602	FAD	P-O3P-PA	-2.74	123.41	132.83
2	A	601	NAI	C4A-C5A-N7A	-2.68	106.61	109.40
2	A	601	NAI	C3D-C2D-C1D	2.54	106.25	101.43
3	A	602	FAD	C10-N1-C2	2.46	121.83	116.90
3	A	602	FAD	C4X-C10-N10	2.36	119.94	116.48
3	A	602	FAD	C4X-C4-N3	2.22	118.83	113.19
3	A	602	FAD	C4'-C3'-C2'	-2.17	108.85	113.36
3	A	602	FAD	O4-C4-C4X	-2.14	120.93	126.60
2	A	601	NAI	C3B-C2B-C1B	2.01	104.00	100.98

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	FAD	O4B-C4B-C5B-O5B
3	A	602	FAD	N10-C1'-C2'-O2'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	602	FAD	N10-C1'-C2'-C3'
3	A	602	FAD	C3'-C4'-C5'-O5'
3	A	602	FAD	O4'-C4'-C5'-O5'
3	A	602	FAD	C5'-O5'-P-O1P
3	A	602	FAD	C5'-O5'-P-O2P
3	A	602	FAD	PA-O3P-P-O5'
2	A	601	NAI	O4D-C1D-N1N-C2N
2	A	601	NAI	C3D-C4D-C5D-O5D
3	A	602	FAD	C3B-C4B-C5B-O5B
3	A	602	FAD	O3'-C3'-C4'-O4'
2	A	601	NAI	O4D-C4D-C5D-O5D
3	A	602	FAD	C2'-C3'-C4'-O4'
3	A	602	FAD	C2'-C3'-C4'-C5'
3	A	602	FAD	O3'-C3'-C4'-C5'
2	A	601	NAI	PA-O3-PN-O5D
2	A	601	NAI	PN-O3-PA-O2A
3	A	602	FAD	P-O3P-PA-O1A
2	A	601	NAI	C2N-C3N-C7N-N7N
2	A	601	NAI	C2N-C3N-C7N-O7N
2	A	601	NAI	PN-O3-PA-O1A
3	A	602	FAD	C5'-O5'-P-O3P
2	A	601	NAI	O4B-C4B-C5B-O5B
3	A	602	FAD	P-O3P-PA-O2A


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	FAD	1	0
2	A	601	NAI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	520/548 (94%)	0.57	40 (7%)	21 20	25, 41, 66, 110	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	325	ALA	4.3
1	A	328	PRO	4.0
1	A	324	TYR	3.7
1	A	482	THR	3.4
1	A	484	PHE	3.4
1	A	504	TYR	3.3
1	A	488	LYS	3.3
1	A	487	ALA	3.2
1	A	2	THR	3.2
1	A	326	CYS	3.1
1	A	502	THR	3.1
1	A	458	ILE	3.1
1	A	459	ASN	3.1
1	A	226	ALA	3.0
1	A	323	LEU	2.9
1	A	4	LYS	2.9
1	A	3	GLN	2.9
1	A	529	LEU	2.7
1	A	394	GLY	2.7
1	A	356	ILE	2.7
1	A	327	ARG	2.7
1	A	501	ASN	2.6
1	A	485	ALA	2.6
1	A	350	ALA	2.5
1	A	114	TYR	2.4
1	A	149	TRP	2.3
1	A	456	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	151	ILE	2.3
1	A	121	PHE	2.2
1	A	398	ILE	2.2
1	A	478	ILE	2.2
1	A	483	LEU	2.2
1	A	530	LEU	2.1
1	A	453	ALA	2.1
1	A	467	ALA	2.1
1	A	400	ILE	2.1
1	A	379	GLY	2.1
1	A	257	SER	2.1
1	A	533	SER	2.1
1	A	57	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAI	A	601	44/44	0.87	0.12	39,51,64,72	0
3	FAD	A	602	53/53	0.93	0.10	27,37,45,48	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAI A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.