



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 31, 2025 – 08:26 PM JST

PDB ID : 4Z1I / pdb\_00004z1i  
Title : Crystal structure of human Trap1 with AMPPNP  
Authors : Lee, C.; Park, H.K.; Ryu, J.H.; Kang, B.H.  
Deposited on : 2015-03-27  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.42


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1085 (3.32-3.28)
Clashscore	180529	1128 (3.32-3.28)
Ramachandran outliers	177936	1125 (3.32-3.28)
Sidechain outliers	177891	1124 (3.32-3.28)
RSRZ outliers	164620	1085 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 45%, green 54%, grey 27%);"></div> <div> <span>%</span> <span style="margin-left: 10px;">45%</span> <span style="margin-left: 10px;">25%</span> <span style="margin-left: 10px;">••</span> <span style="margin-left: 10px;">27%</span> </div> </div>
1	B	502	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 65%, green 34%, grey 6%);"></div> <div> <span>%</span> <span style="margin-left: 10px;">65%</span> <span style="margin-left: 10px;">27%</span> <span style="margin-left: 10px;">••</span> <span style="margin-left: 10px;">6%</span> </div> </div>
1	C	502	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 61%, green 38%, grey 6%);"></div> <div> <span>%</span> <span style="margin-left: 10px;">61%</span> <span style="margin-left: 10px;">29%</span> <span style="margin-left: 10px;">••</span> <span style="margin-left: 10px;">6%</span> </div> </div>
1	D	502	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 59%, green 40%, grey 11%);"></div> <div> <span>%</span> <span style="margin-left: 10px;">59%</span> <span style="margin-left: 10px;">27%</span> <span style="margin-left: 10px;">••</span> <span style="margin-left: 10px;">11%</span> </div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 14236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock protein 75 kDa, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	365	Total	C	N	O	S	0	0	0
			2911	1855	499	547	10			
1	B	472	Total	C	N	O	S	0	0	0
			3798	2417	648	719	14			
1	C	471	Total	C	N	O	S	0	0	0
			3787	2409	647	718	13			
1	D	449	Total	C	N	O	S	0	0	0
			3612	2303	614	682	13			

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	115.55Å 115.55Å 339.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.82 – 3.30 37.82 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (37.82-3.30) 92.5 (37.82-3.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.97 (at 3.32Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.220 , 0.285 0.229 , 0.289	Depositor DCC
$R_{free}$ test set	1892 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.5	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 66.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.067 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	14236	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.42	0/2963	0.80	9/3987 (0.2%)
1	B	0.39	0/3871	0.75	6/5213 (0.1%)
1	C	0.52	6/3859 (0.2%)	0.85	16/5198 (0.3%)
1	D	0.48	3/3677 (0.1%)	0.82	20/4948 (0.4%)
All	All	0.46	9/14370 (0.1%)	0.81	51/19346 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	3
1	C	0	4
1	D	0	1
All	All	0	12

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	498	TYR	CD1-CE1	-12.09	1.21	1.39
1	C	499	TYR	CD1-CE1	9.66	1.53	1.39
1	C	498	TYR	CD2-CE2	-8.35	1.26	1.39
1	C	498	TYR	CE2-CZ	-7.96	1.28	1.38
1	C	498	TYR	CB-CG	-7.18	1.40	1.51

The worst 5 of 51 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	191	ALA	N-CA-C	12.19	143.91	111.00
1	B	414	LEU	CA-CB-CG	11.32	141.33	115.30
1	B	439	LYS	CD-CE-NZ	10.06	134.85	111.70
1	B	411	ILE	CG1-CB-CG2	-9.64	90.18	111.40
1	C	192	GLU	N-CA-CB	-9.34	93.79	110.60

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	407	GLU	Peptide
1	A	434	ALA	Peptide
1	A	449	ARG	Peptide
1	A	452	ILE	Peptide
1	B	411	ILE	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2911	0	2936	147	0
1	B	3798	0	3799	145	0
1	C	3787	0	3793	175	0
1	D	3612	0	3625	170	0
2	A	31	0	13	1	0
2	B	31	0	13	0	0
2	C	31	0	13	1	0
2	D	31	0	13	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	14236	0	14205	619	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 619 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:498:TYR:CZ	1:C:549:SER:HA	1.97	0.99
1:B:228:GLY:HA3	1:B:247:VAL:HG21	1.46	0.98
1:C:497:ILE:HG13	1:C:545:LYS:HD2	1.48	0.95
1:D:408:SER:O	1:D:410:LEU:N	1.99	0.94
1:B:411:ILE:HG22	1:B:415:ARG:NH1	1.82	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/502 (71%)	310 (87%)	35 (10%)	12 (3%)	3	19
1	B	466/502 (93%)	433 (93%)	30 (6%)	3 (1%)	22	53
1	C	465/502 (93%)	442 (95%)	19 (4%)	4 (1%)	14	44
1	D	437/502 (87%)	412 (94%)	21 (5%)	4 (1%)	14	44
All	All	1725/2008 (86%)	1597 (93%)	105 (6%)	23 (1%)	10	36

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	PRO
1	A	293	ASN
1	A	336	PRO
1	A	435	GLU
1	A	437	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/443 (72%)	308 (97%)	10 (3%)	35	61
1	B	415/443 (94%)	409 (99%)	6 (1%)	62	78
1	C	414/443 (94%)	405 (98%)	9 (2%)	47	69
1	D	398/443 (90%)	391 (98%)	7 (2%)	54	74
All	All	1545/1772 (87%)	1513 (98%)	32 (2%)	48	70

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	395	ASP
1	D	408	SER
1	B	412	ARG
1	B	408	SER
1	D	412	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	399	ASN
1	C	322	HIS
1	D	136	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ANP	C	801	1,3	29,33,33	1.19	4 (13%)	31,52,52	1.07	3 (9%)
2	ANP	D	801	1,3	29,33,33	2.04	4 (13%)	31,52,52	2.73	7 (22%)
2	ANP	B	801	1,3	29,33,33	1.11	4 (13%)	31,52,52	1.33	3 (9%)
2	ANP	A	801	3	29,33,33	2.13	6 (20%)	31,52,52	1.05	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	C	801	1,3	-	9/14/38/38	0/3/3/3
2	ANP	D	801	1,3	-	10/14/38/38	0/3/3/3
2	ANP	B	801	1,3	-	3/14/38/38	0/3/3/3
2	ANP	A	801	3	-	6/14/38/38	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	ANP	PG-O1G	7.58	1.58	1.46
2	D	801	ANP	PG-O1G	7.52	1.58	1.46
2	A	801	ANP	PB-O1B	6.36	1.56	1.46
2	D	801	ANP	PB-O1B	5.56	1.55	1.46
2	C	801	ANP	PB-O3A	-3.21	1.55	1.59

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	801	ANP	O1G-PG-N3B	-12.01	94.08	111.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	801	ANP	PB-O3A-PA	-5.19	114.34	132.62
2	B	801	ANP	PB-O3A-PA	-4.75	115.89	132.62
2	D	801	ANP	O1B-PB-N3B	-4.66	104.91	111.77
2	C	801	ANP	PB-O3A-PA	-4.04	118.40	132.62

There are no chirality outliers.

5 of 28 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	ANP	PB-N3B-PG-O1G
2	A	801	ANP	PG-N3B-PB-O1B
2	A	801	ANP	PA-O3A-PB-O1B
2	A	801	ANP	PA-O3A-PB-O2B
2	B	801	ANP	PB-N3B-PG-O1G


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	801	ANP	1	0
2	D	801	ANP	3	0
2	A	801	ANP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	365/502 (72%)	-0.03	5 (1%) 73 60	39, 75, 149, 189	1 (0%)
1	B	472/502 (94%)	-0.17	5 (1%) 77 66	35, 67, 113, 170	1 (0%)
1	C	471/502 (93%)	-0.06	5 (1%) 77 66	36, 81, 144, 195	1 (0%)
1	D	449/502 (89%)	-0.02	7 (1%) 70 57	35, 78, 144, 193	0
All	All	1757/2008 (87%)	-0.07	22 (1%) 74 61	35, 74, 141, 195	3 (0%)

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	TYR	4.2
1	C	191	ALA	4.0
1	C	193	ALA	3.2
1	C	192	GLU	2.9
1	D	411	ILE	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	C	802	1/1	0.86	0.09	98,98,98,98	0
3	MG	D	802	1/1	0.89	0.13	215,215,215,215	0
3	MG	A	802	1/1	0.90	0.12	42,42,42,42	0
3	MG	B	802	1/1	0.92	0.08	36,36,36,36	0
2	ANP	D	801	31/31	0.94	0.09	53,59,87,108	0
2	ANP	C	801	31/31	0.96	0.08	42,48,74,125	0
2	ANP	A	801	31/31	0.96	0.07	37,40,53,99	0
2	ANP	B	801	31/31	0.97	0.07	13,51,55,129	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around ANP C 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ANP A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.