wwPDB EM Validation Summary Report

Oct 25, 2026 – 11:33 AM EDT

PDB ID	:	9ZVQ / pdb_00009zvq
EMDB ID	:	EMD-74888
Title	:	EBV gB trimer with "deliDoP" mutations and I5350A2hN (prefusion conformation) - I5350A2hN local refinement
Authors	:	McCallum, M.; Veesler, D.; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on	:	2025-12-30
Resolution	:	2.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10327 (2.20 - 3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	907	25% 72%
1	B	907	25% 72%
1	C	907	25% 72%

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 5568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EBV gB trimer with "deliDoP" mutations and I5350A2hN.


Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	254	Total	C	N	O	S	0	0
			1856	1203	320	326	7		
1	B	254	Total	C	N	O	S	0	0
			1856	1203	320	326	7		
1	C	254	Total	C	N	O	S	0	0
			1856	1203	320	326	7		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: EBV gB trimer with "deliDoP" mutations and I5350A2hN

Chain A:




• Molecule 1: EBV gB trimer with "deliDoP" mutations and I5350A2hN


Chain B:



• Molecule 1: EBV gB trimer with "deliDoP" mutations and I5350A2hN

Chain C:




ALA PRO GLY SER THR THR GLN THR THR ARG THR ARG THR THR VAL ASN CYS LEU ILE THR ASP MET MET ALA LYS SER ASN SER PRO PHE ASP PHE PHE VAL THR THR THR GLY GLN THR VAL GLU MET SER PRO PHE TYR ASP GLY LYS ASN LYS GLU THR PHE HIS GLU ARG ALA ASP
SER PHE HIS VAL ARG THR ASN TYR LYS ILE VAL ASP TYR ASP ASN ARG GLY THR ASN PRO GLN GLY GLU ARG ARG ALA PHE LEU ASP LYS GLY THR TYR THR LEU SER TRP LYS LEU GLU ASN ARG THR PRO TYR CYS PRO LEU GLN HIS TRP GLN THR PHE ASP SER THR ILE ALA THR
GLU THR GLY LYS SER ILE HIS PHE VAL THR ASP GLU GLY THR SER SER PHE VAL THR ASN THR THR VAL GLY ILE GLU LEU PRO ASP ALA PHE LYS CYS ILE GLU GLU GLN VAL ASN LYS THR MET HIS GLU LYS TYR GLU ALA VAL GLN ILE GLN ASP ARG TYR THR LYS GLY GLN GLU ALA ILE
THR TYR PHE ILE THR SER GLY GLY LEU LEU LEU ALA TRP LEU PRO LEU THR PRO ARG SER LEU ALA THR VAL LYS ASN LEU THR GLU LEU THR THR THR GLY SER PRO PRO GLY SER PRO PRO LEU ASN ASN PRO ALA THR VAL GLN ILE GLN PHE ALA TYR ASP SER LEU ARG GLN ILE
ASN ARG MET LEU GLY ASP LEU ALA ARG GLN LYS ARG GLN ASN MET VAL LEU ARG GLU LEU THR LYS ILE ASN PRO THR THR VAL MET SER SER ILE TYR GLY LYS ALA VAL ALA ALA LYS ARG LEU GLY ASP VAL ILE GLN PHE ALA TYR VAL SER CYS CYS VAL PRO VAL ASN
GLN ALA THR VAL THR LEU ARG LYS SER MET ARG VAL PRO GLY SER GLU THR MET CYS TYR SER ARG PRO LEU VAL SER PHE SER PHE ILE ASN ASP THR LYS THR TYR GLU GLY GLN LEU GLY THR ASP ASN GLU ILE PHE LEU THR LYS LYS MET THR GLU VAL CYS GLN ALA THR SER
GLN TYR TYR PHE GLN SER GLY ASN GLU ILE HIS VAL TYR ASN ASP TYR HIS HIS PHE LYS THR ILE GLU LEU ASP GLY ILE ALA THR LEU GLN THR PHE ILE SER LEU ASN THR SER LEU ILE CYS ASN ILE ASP PHE ALA SER LEU GLU LEU TYR SER ARG ASP GLU GLN ARG ALA SER
ASN VAL PHE ASP LEU GLU GLY ILE PHE ARG GLU LEU PRO PHE ILE ALA GLN ASN ILE ALA ALA ILE GLU LYS ARG ILE ASN GLU ILE GLU ARG ARG ILE SER GLY SER GLY SER GLY GLY SER GLY HIS HIS HIS HIS HIS HIS HIS HIS

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	155157	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.560	Depositor
Minimum map value	-0.543	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.017	Depositor
Recommended contour level	0.4	Depositor
Map size (Å)	424.448, 424.448, 424.448	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.829, 0.829, 0.829	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.11	0/1887	0.27	0/2552
1	B	0.11	0/1887	0.27	0/2552
1	C	0.11	0/1887	0.27	0/2552
All	All	0.11	0/5661	0.27	0/7656

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1856	0	1894	13	0
1	B	1856	0	1894	13	0
1	C	1856	0	1894	12	0
All	All	5568	0	5682	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:LEU:HD12	1:A:326:LEU:HD22	1.81	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100:LEU:HD12	1:B:326:LEU:HD22	1.81	0.61
1:C:100:LEU:HD12	1:C:326:LEU:HD22	1.81	0.61
1:C:173:VAL:O	1:C:192:SER:OG	2.21	0.57
1:A:173:VAL:O	1:A:192:SER:OG	2.21	0.56

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/907 (28%)	249 (99%)	3 (1%)	0	100	100
1	B	252/907 (28%)	249 (99%)	3 (1%)	0	100	100
1	C	252/907 (28%)	249 (99%)	3 (1%)	0	100	100
All	All	756/2721 (28%)	747 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/789 (23%)	181 (100%)	1 (0%)	81	92
1	B	182/789 (23%)	181 (100%)	1 (0%)	81	92
1	C	182/789 (23%)	181 (100%)	1 (0%)	81	92

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	546/2367 (23%)	543 (100%)	3 (0%)	78 92

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	LYS
1	B	204	LYS
1	C	204	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-74888. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 256



Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256




Z Index: 256

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 240




Y Index: 248




Z Index: 316


6.3.2 Raw map



X Index: 240



Y Index: 248




Z Index: 316


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 28 nm^3; this corresponds to an approximate mass of 26 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.370 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.370 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.70	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.67	6.25	3.74

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.67 differs from the reported value 2.7 by more than 10 %


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-74888 and PDB model 9ZVQ. Per-residue inclusion information can be found in section 3 on page 4.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.4).

9.4 Atom inclusion (i)



At the recommended contour level, 89% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.4) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.7770	0.5590
A	0.7770	0.5580
B	0.7750	0.5600
C	0.7800	0.5590