



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 14, 2026 – 10:08 AM EDT

PDB ID : 9ZX5 / pdb_00009zx5
Title : Crystal structure of the N4BP2 polynucleotide kinase domain complexed with ATP or ADP, Mg, and the trinucleotide substrate UGA.
Authors : Wang, H.; Stanley, R.E.
Deposited on : 2026-01-03
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)
RNA backbone	3983	1027 (2.66-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	% 79% 5% 16%
1	B	182	5% 79% 5% 16%
1	C	182	% 79% 7% 15%
2	D	3	67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	3	 33% 67%
2	F	3	 67% 33%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8129 atoms, of which 3849 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEDD4-binding protein 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	153	Total	C	H	N	O	S	0	0	0
			2476	809	1229	211	224	3			
1	B	153	Total	C	H	N	O	S	0	0	0
			2490	811	1238	214	224	3			
1	C	155	Total	C	H	N	O	S	0	0	0
			2544	825	1271	218	227	3			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	414	GLY	-	expression tag	UNP Q86UW6
A	415	SER	-	expression tag	UNP Q86UW6
A	416	HIS	-	expression tag	UNP Q86UW6
A	417	MET	-	expression tag	UNP Q86UW6
A	418	ALA	-	expression tag	UNP Q86UW6
A	419	THR	-	expression tag	UNP Q86UW6
A	420	ASP	-	expression tag	UNP Q86UW6
A	421	TYR	-	expression tag	UNP Q86UW6
A	422	LYS	-	expression tag	UNP Q86UW6
A	423	ASP	-	expression tag	UNP Q86UW6
A	424	ASP	-	expression tag	UNP Q86UW6
A	425	ASP	-	expression tag	UNP Q86UW6
A	426	ASP	-	expression tag	UNP Q86UW6
A	427	LYS	-	expression tag	UNP Q86UW6
B	414	GLY	-	expression tag	UNP Q86UW6
B	415	SER	-	expression tag	UNP Q86UW6
B	416	HIS	-	expression tag	UNP Q86UW6
B	417	MET	-	expression tag	UNP Q86UW6
B	418	ALA	-	expression tag	UNP Q86UW6
B	419	THR	-	expression tag	UNP Q86UW6
B	420	ASP	-	expression tag	UNP Q86UW6
B	421	TYR	-	expression tag	UNP Q86UW6
B	422	LYS	-	expression tag	UNP Q86UW6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	423	ASP	-	expression tag	UNP Q86UW6
B	424	ASP	-	expression tag	UNP Q86UW6
B	425	ASP	-	expression tag	UNP Q86UW6
B	426	ASP	-	expression tag	UNP Q86UW6
B	427	LYS	-	expression tag	UNP Q86UW6
C	414	GLY	-	expression tag	UNP Q86UW6
C	415	SER	-	expression tag	UNP Q86UW6
C	416	HIS	-	expression tag	UNP Q86UW6
C	417	MET	-	expression tag	UNP Q86UW6
C	418	ALA	-	expression tag	UNP Q86UW6
C	419	THR	-	expression tag	UNP Q86UW6
C	420	ASP	-	expression tag	UNP Q86UW6
C	421	TYR	-	expression tag	UNP Q86UW6
C	422	LYS	-	expression tag	UNP Q86UW6
C	423	ASP	-	expression tag	UNP Q86UW6
C	424	ASP	-	expression tag	UNP Q86UW6
C	425	ASP	-	expression tag	UNP Q86UW6
C	426	ASP	-	expression tag	UNP Q86UW6
C	427	LYS	-	expression tag	UNP Q86UW6

- Molecule 2 is a RNA chain called RNA (5'-r(UpGpA)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	3	Total	C	H	N	O	P	0	0	0
			66	19	22	7	16	2			
2	E	3	Total	C	H	N	O	P	0	0	0
			66	19	22	7	16	2			
2	F	3	Total	C	H	N	O	P	0	0	0
			66	19	22	7	16	2			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃)


ATP

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 43	C 10	H 12	N 5	O 13	P 3	0	0
4	B	1	Total 43	C 10	H 12	N 5	O 13	P 3	0	0
4	C	1	Total 56	C 11	H 15	N 5	O 20	P 5	0	1

-

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	H	O	0	0
			10	2	6	2		



- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	65	Total	O	0	0
			65	65		
7	B	67	Total	O	0	0
			67	67		
7	C	110	Total	O	0	2
			110	110		
7	D	5	Total	O	0	0
			5	5		
7	E	4	Total	O	0	0
			4	4		
7	F	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NEDD4-binding protein 2





- Molecule 1: NEDD4-binding protein 2



GLU

- Molecule 1: NEDD4-binding protein 2



- Molecule 2: RNA (5'-r(UpGpA)-3')



U1 G2 A3

- Molecule 2: RNA (5'-r(UpGpA)-3')



U1 G2 A3

- Molecule 2: RNA (5'-r(UpGpA)-3')

Chain F:



Residue	Color
11	Green
62	Yellow
13	Green

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	103.98Å 106.20Å 112.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.13 – 2.35 33.13 – 2.35	Depositor EDS
% Data completeness (in resolution range)	96.7 (33.13-2.35) 96.6 (33.13-2.35)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 2.22Å)	Xtriage
Refinement program	PHENIX 2.0_5936	Depositor
R, R_{free}	0.205 , 0.243 0.202 , 0.240	Depositor DCC
R_{free} test set	1360 reflections (4.34%)	wwPDB-VP
Wilson B-factor (Å ²)	42.4	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.005 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8129	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, ATP, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/1281	0.31	0/1736
1	B	0.12	0/1286	0.31	0/1742
1	C	0.12	0/1307	0.32	0/1768
2	D	0.11	0/48	0.31	0/74
2	E	0.13	0/48	0.28	0/74
2	F	0.12	0/48	0.23	0/74
All	All	0.12	0/4018	0.31	0/5468

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1247	1229	1229	5	0
1	B	1252	1238	1238	5	0
1	C	1273	1271	1271	7	0
2	D	44	22	22	0	0
2	E	44	22	22	1	0
2	F	44	22	22	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	31	12	12	0	0
4	B	31	12	12	0	0
4	C	41	15	6	1	0
5	B	10	0	0	0	0
6	C	4	6	6	0	0
7	A	65	0	0	0	0
7	B	67	0	0	0	0
7	C	110	0	0	0	0
7	D	5	0	0	0	0
7	E	4	0	0	0	0
7	F	5	0	0	0	0
All	All	4280	3849	3840	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:552:GLU:HG3	1:B:556:ARG:HE	1.66	0.60
1:C:456:LEU:HD21	1:C:540:ARG:HD2	1.92	0.51
1:C:490:GLY:O	1:C:494:GLU:HG2	2.11	0.50
1:B:565:LYS:HE3	2:E:1:U:O2	2.12	0.49
1:A:491:GLU:HA	1:A:494:GLU:HG2	1.95	0.48

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/182 (83%)	148 (98%)	3 (2%)	0	100	100
1	B	151/182 (83%)	146 (97%)	5 (3%)	0	100	100
1	C	153/182 (84%)	150 (98%)	3 (2%)	0	100	100
All	All	455/546 (83%)	444 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/165 (81%)	132 (98%)	2 (2%)	57	72
1	B	135/165 (82%)	135 (100%)	0	100	100
1	C	138/165 (84%)	138 (100%)	0	100	100
All	All	407/495 (82%)	405 (100%)	2 (0%)	81	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	438	VAL
1	A	540	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	533	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	1/3 (33%)	1 (100%)	0
2	E	1/3 (33%)	1 (100%)	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	F	1/3 (33%)	1 (100%)	0
All	All	3/9 (33%)	3 (100%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	2	G
2	E	2	G
2	F	2	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ATP	A	2002	3	32,33,33	0.67	0	48,52,52	0.63	1 (2%)
4	ATP	C	2002[B]	3	32,29,33	0.76	0	48,45,52	0.64	1 (2%)
4	ATP	C	2002[A]	3	32,33,33	0.71	0	48,52,52	0.57	0
4	ATP	B	2002	3	32,33,33	0.74	0	48,52,52	0.50	0
6	EDO	C	2003	-	3,3,3	0.23	0	2,2,2	0.38	0
5	SO4	B	2004	-	4,4,4	0.68	0	6,6,6	0.09	0
5	SO4	B	2003	-	4,4,4	0.67	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	2002	3	-	2/22/38/38	0/3/3/3
4	ATP	C	2002[B]	3	-	2/22/32/38	0/3/3/3
4	ATP	C	2002[A]	3	-	1/22/38/38	0/3/3/3
6	EDO	C	2003	-	-	0/1/1/1	-
4	ATP	B	2002	3	-	2/22/38/38	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2002	ATP	O3G-PG-O2G	2.29	116.39	107.80
4	C	2002[B]	ATP	O2G-PG-O3B	2.11	111.70	104.64

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2002	ATP	PA-O3A-PB-O1B
4	B	2002	ATP	PA-O3A-PB-O1B
4	B	2002	ATP	PA-O3A-PB-O2B
4	C	2002[B]	ATP	PA-O3A-PB-O2B
4	A	2002	ATP	PA-O3A-PB-O2B


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	2002[A]	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/182 (84%)	0.07	2 (1%) 75 79	32, 48, 76, 98	0
1	B	153/182 (84%)	0.35	10 (6%) 25 28	33, 56, 86, 104	0
1	C	155/182 (85%)	-0.10	2 (1%) 75 79	28, 43, 68, 90	0
2	D	3/3 (100%)	-0.79	0 100 100	44, 44, 47, 58	0
2	E	3/3 (100%)	-0.40	0 100 100	46, 46, 51, 65	0
2	F	3/3 (100%)	-0.97	0 100 100	35, 35, 37, 46	0
All	All	470/555 (84%)	0.09	14 (2%) 52 59	28, 49, 80, 104	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	437	TYR	4.6
1	B	478	ILE	4.0
1	B	436	SER	3.0
1	B	479	ASN	2.7
1	A	588	PRO	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	C	2003	4/4	0.75	0.20	52,63,71,85	0
5	SO4	B	2004	5/5	0.93	0.08	43,51,55,61	0
4	ATP	C	2002[B]	27/31	0.97	0.07	25,31,40,42	13
5	SO4	B	2003	5/5	0.97	0.07	35,37,47,50	0
4	ATP	A	2002	31/31	0.97	0.07	27,39,50,52	0
4	ATP	C	2002[A]	31/31	0.97	0.07	24,31,38,42	17
4	ATP	B	2002	31/31	0.98	0.05	27,33,41,41	0
3	MG	B	2001	1/1	0.99	0.04	30,30,30,30	0
3	MG	C	2001	1/1	0.99	0.03	26,26,26,26	0
3	MG	A	2001	1/1	0.99	0.02	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around EDO C 2003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP C 2002 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP A 2002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP C 2002 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP B 2002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG B 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around MG C 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.